arXiv:q-alg/9606005v1 5 Jun 1996

SOLUTIONS OF THE ELLIPTIC QKZB EQUATIONS AND BETHE ANSATZ I

G. FELDER*,[1], V. TARASOV**,[2] AND A. VARCHENKO*,[3]

Department of Mathematics, University of North Carolina at Chapel Hill,
Chapel Hill, NC 27599-3250, USA
E-mail addresses: felder@math.unc.edu, varchenko@math.unc.edu

***Laboratoire de Physique Théorique ENSLAPP, École Normale Supérieure de Lyon,*
46, Allée d'Italie, 69364 Lyon Cedex 07, France
E-mail address: vtarasov@enslapp.ens-lyon.fr

June 1996

Dedicated to V. I. Arnold on his sixtieth birthday

1. Introduction

In this paper we give integral representations for solutions of the system of elliptic quantum Knizhnik–Zamolodchikov–Bernard (qKZB) difference equations in the case of sl_2. The qKZB equations [F] are a quantum deformation of the KZB differential equations obeyed by correlation functions of the Wess–Zumino–Witten model on tori. They have the form

$$\Psi(z_1,\dots,z_j+p,\dots,z_n)=K_j(z_1,\dots,z_n;\tau,\eta,p)\Psi(z_1,\dots,z_n).$$

The unknown function Ψ takes values in a space of vector valued functions of a complex variable λ, and the K_j are difference operators in λ. The parameters of this system of equations are τ (the period of the elliptic curve), η ("Planck's constant"), p (the step) and n "highest weights" $\Lambda_1,\dots,\Lambda_n\in\mathbb{C}$. The operators K_j are expressed in terms of R-matrices of the elliptic quantum group $E_{\tau,\eta}(sl_2)$.

In the trigonometric limit $\tau\to i\infty$, the qKZB equations reduce to the trigonometric qKZ equations [FR] obeyed by correlation functions of statistical models and form factors of integrable quantum field theories in 1+1 dimensions.

The KZB equations can be obtained in the semiclassical limit: $\eta\to0$, $p\to0$, p/η finite.

When the step p of the qKZB equations goes to zero (with the other parameters fixed) our construction gives common eigenfunctions of the n commuting operators $K_j(z_1,\dots,z_n;\tau,\eta,0)$ in the form of the Bethe ansatz. These difference operators are closely related to the transfer matrices of IRF models of statistical mechanics.

Our results follow from the main theme of this paper: a geometric construction of tensor products of evaluation Verma modules over the elliptic quantum group $E_{\tau,\eta}(sl_2)$. In particular, we obtain some formulae given in [FV1] for the action of generators on these modules.

---

[1] Supported in part by NSF grant DMS-9400841
Address after July 1, 1996: D-Math, ETH Zentrum, CH-8092 Zürich, Switzerland.

[2] Supported in part by MAE–MICECO–CNRS Fellowship.

[3] Supported in part by NSF grant DMS-9501290.

The results of this paper are thus parallel to the results on solutions of the rational and trigonometric qKZB equations of [V, TV2], which are based on the representation theory of the Yangian $Y(sl_2)$ and the affine quantum universal enveloping algebra $U_q(\widehat{sl_2})$, respectively.

The paper is organized as follows: we begin by introducing the notion of R-matrices and the qKZB equations in Section 2. The geometric construction of R-matrices is given in Section 3. At the end of that section we show how to obtain representations of $E_{\tau,\eta}(sl_2)$ in this way, and give some explicit formulae for R-matrix elements.

The main applications of the geometric construction are contained in Sections 4–7. The Bethe ansatz eigenfunctions of the commuting system of difference operators are given in Section 4, Theorem 13. The proof of completeness of Bloch eigenfunctions obtained by the Bethe ansatz in the case of rational η is given for generic parameters in Section 5. In Section 6, we compare our results with the algebraic Bethe ansatz of [FV2]. In particular, we show, using the results of Section 5 that the algebraic Bethe ansatz gives a basis of eigenvectors in some cases. Our results on integral representation for solutions of the qKZB equations are stated in Theorem 22 and Theorem 24, Section 7. In Section 8 we introduce the technique of iterated residues, the main calculational tool of the paper, and complete the proofs of our results.

Some of our results seem to be related to the results in the recent preprint [T], in which integral solutions of a system of difference equations associated to Sklyanin's algebra are constructed.

In the next paper of this series, we will discuss integration cycles, and compute the monodromy of the qKZB equations. It turns out that the monodromy is described in terms of R-matrices of the elliptic quantum group $E_{p,\eta}(sl_2)$ and that the elliptic quantum groups associated to the elliptic curves $\mathbb{C}/(\mathbb{Z}+\tau\mathbb{Z})$ and $\mathbb{C}/(\mathbb{Z}+p\mathbb{Z})$ play a symmetric role in the story.

2. R-MATRICES, qKZB EQUATIONS AND COMMUTING DIFFERENCE OPERATORS

2.1. R-matrices.

The qKZB equations are given in terms of R-matrices of elliptic quantum groups. In the sl_2 case, these R-matrices have the following properties. Let $\mathfrak{h}=\mathbb{C}h$ be a one-dimensional Lie algebra with generator h. For each $\Lambda\in\mathbb{C}$ consider the $\mathfrak{h}$-module $V_\Lambda=\oplus_{j=0}^{\infty}\mathbb{C}e_j$, with $he_j=(\Lambda-2j)e_j$. For each pair Λ_1, Λ_2 of complex numbers we have a meromorphic function, called the R-matrix, $R_{\Lambda_1,\Lambda_2}(z,\lambda)$ of two complex variables, with values in $\mathrm{End}(V_{\Lambda_1}\otimes V_{\Lambda_2})$.

The main properties of the R-matrices are

I. Zero weight property: for any Λ_i, z,λ, $[R_{\Lambda_1,\Lambda_2}(z,\lambda),h^{(1)}+h^{(2)}]=0$.

II. For any $\Lambda_1,\Lambda_2,\Lambda_3$, the dynamical Yang–Baxter equation

$$\begin{aligned}R_{\Lambda_1,\Lambda_2}(z,\lambda-2\eta h^{(3)})^{(12)}R_{\Lambda_1,\Lambda_3}(z+w,\lambda)^{(13)}R_{\Lambda_2,\Lambda_3}(w,\lambda-2\eta h^{(1)})^{(23)}\\ =R_{\Lambda_2,\Lambda_3}(w,\lambda)^{(23)}R_{\Lambda_1,\Lambda_3}(z+w,\lambda-2\eta h^{(2)})^{(13)}R_{\Lambda_1,\Lambda_2}(z,\lambda)^{(12)},\end{aligned}$$

holds in $\mathrm{End}(V_{\Lambda_1}\otimes V_{\Lambda_2}\otimes V_{\Lambda_3})$ for all z,w,λ.

III. For all Λ_1, Λ_2, z,λ, $R_{\Lambda_1,\Lambda_2}(z,\lambda)^{(12)}R_{\Lambda_2,\Lambda_1}(-z,\lambda)^{(21)}=\mathrm{Id}$. This property is called "unitarity".

We use the following notation: if $X\in\mathrm{End}(V_i)$ we denote by $X^{(i)}\in\mathrm{End}(V_1\otimes\cdots\otimes V_n)$ the operator $\cdots\otimes\mathrm{Id}\otimes X\otimes\mathrm{Id}\otimes\cdots$, acting non-trivially on the ith factor of a tensor product of vector spaces, and if $X=\sum X_k\otimes Y_k\in\mathrm{End}(V_i\otimes V_j)$ we set $X^{(ij)}=\sum X_k^{(i)}Y_k^{(j)}$. If $X(\mu_1,\dots,\mu_n)$ is a function with values in $\mathrm{End}(V_1\otimes\cdots\otimes V_n)$, then $X(h^{(1)},\dots,h^{(n)})v=X(\mu_1,\dots,\mu_n)v$ if $h^{(i)}v=\mu_iv$, for all $i=1,\dots,n$.

For each τ in the upper half plane and generic $\eta \in \mathbb{C}$ ("Planck's constant") a system of R-matrices $R_{\Lambda_1,\Lambda_2}(z,\lambda)$ obeying I–III was constructed in [FV1]. They are characterized by an intertwining property with respect to the action of the elliptic quantum group $E_{\tau,\eta}(sl_2)$ on tensor products of evaluation Verma modules. In this paper, we give an alternative geometric construction of these R-matrices, see Section 3.

2.2. **qKZB equations.** Fix the parameters τ, η. Fix also n complex numbers $\Lambda_1, \dots, \Lambda_n$ and an additional parameter $p \in \mathbb{C}$. Let $V = V_{\Lambda_1} \otimes \cdots \otimes V_{\Lambda_n}$. The kernel of $h^{(1)} + \cdots + h^{(n)}$ on V is called the zero-weight space and is denoted $V[0]$. More generally, we write $V[\mu]$ for the eigenspace of $\sum h^{(i)}$ with eigenvalue μ. The qKZB equations are difference equations for a function $\Psi(z_1, \dots, z_n, \lambda)$ of n complex variables $z_1, \dots, z_n$ with values in the space of meromorphic functions $\mathrm{Fun}(V[0])$ of a complex variable λ with values in $V[0]$.

The qKZB equations [F] have the form

$$\begin{aligned}\Psi(z_1, \dots, z_j + p, \dots, z_n) = {} & R_{j,j-1}(z_j - z_{j-1} + p) \cdots R_{j,1}(z_j - z_1 + p) \\ & \Gamma_j R_{j,n}(z_j - z_n) \cdots , R_{j,j+1}(z_j - z_{j+1}) \Psi(z_1, \dots, z_n)\end{aligned} \tag{1}$$

Here $R_{k,l}(z)$ is the operator of multiplication by

$$R_{\Lambda_k,\Lambda_l}(z, \lambda - 2\eta \sum_{\substack{j=1 \\ j \neq k}}^{l-1} h^{(j)})^{(k,l)}$$

acting on the kth and lth factor of the tensor product, and Γ_j is the linear difference operator such that $\Gamma_j \Psi(\lambda) = \Psi(\lambda - 2\eta\mu)$ if $h^{(j)}\Psi = \mu\Psi$.

The consistency of these equations follows from I–III. In other words, the qKZB equations may be viewed as the equation of horizontality for a flat discrete connection on a trivial vector bundle with fiber $\mathrm{Fun}(V[0])$ over an open subset of $\mathbb{C}^n$.

2.3. **Commuting difference operators.** A closely related set of difference equations is the eigenvalue problem

$$H_j(z)\psi = \epsilon_j \psi, \qquad j = 1, \dots, n, \qquad \psi \in \mathrm{Fun}(V[0])$$

for the commuting difference operators

$$H_j(z) = R_{j,j-1}(z_j - z_{j-1}) \cdots R_{j,1}(z_j - z_1) \Gamma_j R_{j,n}(z_j - z_n) \cdots , R_{j,j+1}(z_j - z_{j+1}). \tag{2}$$

Here $z = (z_1, \dots, z_n)$ is a fixed generic point in $\mathbb{C}^n$ and ψ is in $\mathrm{Fun}(V[0])$. The fact that the operators $H_j(z)$ commute with each other follows from the flatness of the connection as $p \to 0$.

2.4. **Finite dimensional representations.** If Λ is a nonnegative integer, V_Λ contains the subspace $SV_\Lambda = \oplus_{j=\Lambda+1}^{\infty} \mathbb{C} e_j$ with the property that, for any M, $SV_\Lambda \otimes V_\mathrm{M}$ and $V_\mathrm{M} \otimes SV_\Lambda$ are preserved by the R-matrices $R_{\Lambda,\mathrm{M}}(z,\lambda)$ and $R_{\mathrm{M},\Lambda}(z,\lambda)$, respectively, see [FV1] and Theorem 8. Let $L_\Lambda = V_\Lambda / SV_\Lambda$, $\Lambda \in \mathbb{Z}_{\geqslant 0}$. Then, in particular, for any nonnegative integers Λ and M, $R_{\Lambda,\mathrm{M}}(z,\lambda)$ induces a map, also denoted by $R_{\Lambda,\mathrm{M}}(z,\lambda)$, on the finite dimensional space $L_\Lambda \otimes L_\mathrm{M}$.

The simplest nontrivial case is $\Lambda = \mathrm{M} = 1$. Then $R_{1,1}(z,\lambda)$ is defined on a four-dimensional vector space and coincides with the *fundamental* R-matrix, the matrix of structure constants of the elliptic quantum group $E_{\tau,\eta}(sl_2)$, see Theorem 10.

In any case, if $\Lambda_1, \dots, \Lambda_n$ are nonnegative integers, we can consider the qKZB equations (1) and the eigenvalue problem (2) on functions with values in the zero weight space of $L_{\Lambda_1} \otimes \cdots \otimes L_{\Lambda_n}$.

The results below obtained for the solutions with values in $\otimes_j V_{\Lambda_j}$ immediately extend to this case: let $\pi : \otimes_{j=1}^n V_{\Lambda_j} \to \otimes_{j=1}^n L_{\Lambda_j}$ denote the canonical projection.

Lemma 1. *Let* $\Psi(z_1,\dots,z_n)$ *be a solution of the qKZB equations with values in* $V[0]=V_{\Lambda_1}\otimes\cdots\otimes V_{\Lambda_n}[0]$. *Then* $\pi\circ\Psi(z_1,\dots,z_n)$ *is a solution of the qKZB equations with values in* $L[0]=L_{\Lambda_1}\otimes\cdots\otimes L_{\Lambda_n}[0]$. *Similarly if* $\Psi\in\mathrm{Fun}(V[0])$ *is a common eigenfunction of the operators* $H_j(z)$ *then* $\pi\circ\Psi$ *is a common eigenfunction of the induced operators* $H_j(z)$ *on* $\mathrm{Fun}(L[0])$*, with the same eigenvalues.*

In this lemma, the zero function should also be understood as an eigenfunction, since, certainly, $\pi\circ\Psi$ can vanish.

2.5. **Remarks.**

1. If $\Lambda_1=\cdots=\Lambda_n=\Lambda\in\mathbb{Z}_{\geqslant 0}$, then the commuting operators $H_j(z)$ on $L[0]$ are special values of the *transfer matrix* $T(w;z_1,\dots,z_n)$. This difference operator is defined as follows. Let $P[\mu]\in\mathrm{End}(L_\Lambda)$ be the projection onto the subspace of L_Λ of weight μ: $P[\Lambda-2j]e_k=\delta_{jk}e_k$, $j=0,\dots,\Lambda$. Define the partial trace

$$\mathrm{tr}_{L_\Lambda[\mu]}=(\mathrm{tr}\circ P[\mu])\otimes\mathrm{Id}:\mathrm{End}(\otimes_{j=0}^n L_\Lambda)\simeq\mathrm{End}(L_\Lambda)\otimes\mathrm{End}(\otimes_{j=1}^n L_\Lambda)\to\mathrm{End}(\otimes_{j=1}^n L_\Lambda)$$

Then the transfer matrix $T(w)=T(w;z_1,\dots,z_n)\in\mathrm{End}(\mathrm{Fun}(L[0]))$ is defined by

$$\begin{aligned} T(w)f(\lambda) &= \mathrm{tr}_{L_\Lambda[\mu]}(R_{0,n}(w-z_n)\cdots R_{0,1}(w-z_1))f(\lambda-2\eta\mu),\\ R_{0,j}(w-z_j) &:= R^{(0,j)}_{\Lambda,\Lambda}(w-z_j,\lambda-2\eta\sum_{k=1}^{j-1}h^{(j)}).\end{aligned}$$

These transfer matrices commute on $L[0]$ for different values of w. It can be shown that

$$H_j(z_1,\dots,z_n)=T(z_j;z_1,\dots,z_n).$$

In particular, the operators $H_j(z_1,\dots,z_n)$ commute with the transfer matrices.

2. Below, a geometric construction of R-matrices is given. This construction gives in particular a construction of $R_{\Lambda,\mathrm{M}}$ on $L_\Lambda\otimes L_\mathrm{M}$ for all $\Lambda,\mathrm{M}\in\mathbb{Z}_{\geqslant 0}$. Alternatively, these R-matrices can be computed starting from $R_{1,1}$ by the *fusion procedure*, see [FV1], Section 8.

3. MODULES OVER THE ELLIPTIC QUANTUM GROUP AS FUNCTION SPACES

In this section we realize the spaces dual to tensor products of evaluation Verma modules over $E_{\tau,\eta}(sl_2)$ as spaces of functions. The R-matrices are then constructed geometrically.

Let us fix complex parameters τ, η with $\mathrm{Im}(\tau)>0$, and complex numbers $\Lambda_1,\dots,\Lambda_n$. We set $a_i=\eta\Lambda_i$, $i=1,\dots,n$.

3.1. **A space of symmetric functions.** We first introduce a space of functions with an action of the symmetric group. Recall that the Jacobi theta function

$$\theta(t)=-\sum_{j\in\mathbb{Z}}e^{\pi i(j+\frac12)^2\tau+2\pi i(j+\frac12)(t+\frac12)},\tag{3}$$

has multipliers -1 and $-\exp(-2\pi it-\pi i\tau)$ as $t\to t+1$ and $t\to t+\tau$, respectively. It is an odd entire function whose zeros are simple and lie on the lattice $\mathbb{Z}+\tau\mathbb{Z}$. It has the product formula

$$\theta(t)\;=\;2e^{\pi i\tau/4}\sin(\pi t)\prod_{j=1}^{\infty}(1-q^j)(1-q^je^{2\pi it})(1-q^je^{-2\pi it}),\qquad q=e^{2\pi i\tau}.$$

Definition: For complex numbers $a_1,\dots,a_n$, $z_1,\dots,z_n$, λ let $\tilde F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ be the space of meromorphic functions $f(t_1,\dots,t_m)$ of m complex variables such that

(i) $\prod_{i<j}\theta(t_i-t_j+2\eta)\prod_{i=1}^m\prod_{k=1}^n\theta(t_i-z_k-a_k)f$ is a holomorphic function on $\mathbb{C}^m$.
(ii) f is periodic with period 1 in each of its arguments and

$$f(\cdots,t_j+\tau,\cdots)=e^{-2\pi i(\lambda+4\eta j-2\eta)}f(\cdots,t_j,\cdots),$$

for all $j=1,\dots,m$.

There is an action of the symmetric group on this space of functions that we introduce now.

Lemma 2. *The symmetric group S_m acts on $\tilde F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ so that the transposition of j and $j+1$ acts as*

$$s_jf(t_1,\dots,t_m)=f(t_1,\dots,t_{j+1},t_j,\dots,t_m)\frac{\theta(t_j-t_{j+1}-2\eta)}{\theta(t_j-t_{j+1}+2\eta)}\,.$$

Proof: Denote by $\phi(t_j-t_{j+1})$ the ratio of theta functions in the definition of the action of s_j. The meromorphic function ϕ is 1-periodic and obeys $\phi(t+\tau)=e^{8\pi i\eta}\phi(t)$. Therefore, the action preserves the behavior of functions in $\tilde F^m$ under translations by the lattice. The position of the poles is also preserved by the action as it is easy to check. The relation $s_js_{j+1}s_j=s_{j+1}s_js_{j+1}$ holds automatically and the relation $s_j^2=1$ follows from $\phi(t)\phi(-t)=1$. These are the two relations defining the symmetric group. □

Definition: For any $m\in\mathbb{Z}_{>0}$, let $F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)=\tilde F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)^{S_m}$ be the space of S_m-invariant functions. If $m=0$, we set $F^0_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)=\mathbb{C}$. We denote by Sym the symmetrization operator Sym$=\sum_{s\in S_m}s:\tilde F^m\to F^m$. Also, we set

$$F_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)=\oplus_{m=0}^\infty F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda),$$

and define an $\mathfrak{h}$-module structure on $F_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda))$ by letting h act by

$$h|_{F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)}=(\sum_{i=1}^n\Lambda_i-2m)\mathrm{Id},\qquad a_i=\eta\Lambda_i.$$

Some technical results on this space of functions are proved in Section 8. In particular, it is shown in Lemma 28 that $F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ is a finite dimensional vector space of dimension $\binom{n+m-1}{m}$.

Clearly, $F^m_{a_{\sigma(1)},\dots,a_{\sigma(n)}}(z_{\sigma(1)},\dots,z_{\sigma(n)})=F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ for any permutation $\sigma\in S_n$.

Example: Let $n=1$. Then $F^m_a(z,\lambda)$ is a one-dimensional space spanned by

$$\omega_m(t_1,\dots,t_m,\lambda;z)=\prod_{i<j}\frac{\theta(t_i-t_j)}{\theta(t_i-t_j+2\eta)}\prod_{j=1}^m\frac{\theta(\lambda+2\eta m+t_j-z-a)}{\theta(t_j-z-a)},\tag{4}$$

see Lemma 29, Section 8.

3.2. **Tensor products.**

Proposition 3. *Let $n = n' + n''$, $m = m' + m''$ be nonnegative integers and $a_1, \dots, a_n$, $z_1, \dots, z_n$ be complex numbers. The formula*

$$k(t_1, \dots, t_m) = \frac{1}{m'!m''!}\mathrm{Sym}\Big(f(t_1, \dots, t_{m'})g(t_{m'+1}, \dots, t_m) \prod_{\substack{m'<j\leqslant m\\ 1\leqslant l\leqslant n'}} \frac{\theta(t_j - z_l + a_l)}{\theta(t_j - z_l - a_l)}\Big)$$

correctly defines a linear map $\Phi : f \otimes g \mapsto k = \Phi(f \otimes g)$,

$$\oplus_{m'=0}^{m} F^{m'}_{a_1,\dots,a_{n'}}(z_1, \dots, z_{n'}, \lambda) \otimes F^{m''}_{a_{n'+1},\dots,a_n}(z_{n'+1}, \dots, z_n, \lambda - 2\nu) \\ \to F^{m}_{a_1,\dots,a_n}(z_1, \dots, z_n, \lambda),$$

where $\nu = a_1 + \dots + a_{n'} - 2\eta m'$. For generic values of the parameters z_j, λ, the map Φ is an isomorphism. Moreover, Φ is associative in the sense that, for any three functions f, g, h, $\Phi(\Phi(f \otimes g) \otimes h) = \Phi(f \otimes \Phi(g \otimes h))$, whenever defined.

The only claim that does not follow immediately from the definitions is the claim that Φ is an isomorphism for generic z_i's and λ. The proof of this is deferred to Section 8.

By iterating this construction, we get for all $n \geqslant 1$ a linear map Φ_n, defined recursively by $\Phi_1 = \mathrm{Id}$, $\Phi_n = \Phi(\Phi_{n-1} \otimes \mathrm{Id})$, from

$$\oplus_{m_1+\dots+m_n=m} \otimes_{i=1}^{n} F^{m_i}_{a_i}(z_i, \lambda - 2\eta(\mu_1 + \dots + \mu_{i-1}))$$

to $F^m_{a_1,\dots,a_n}(z_1, \dots, z_n, \lambda)$, with $\mu_j = a_j/\eta - 2m_j$, $j = 1, \dots, n$. Let $V^*_\Lambda = \oplus_{j=0}^{\infty}\mathbb{C}e^*_j$ be the restricted dual of the module $V_\Lambda = \oplus_{j=0}^{\infty}\mathbb{C}e_j$. It is spanned by the basis (e^*_j) dual to the basis (e_j). We let $\mathfrak{h}$ act on V^*_Λ by $he^*_j = (\Lambda - 2j)e^*_j$. Then the map that sends e^*_j to ω_j (see (4)) defines an isomorphism of $\mathfrak{h}$-modules

$$\omega(z, \lambda) : V^*_\Lambda \to F_a(z, \lambda), \qquad a = \eta\Lambda.$$

By composing this with the maps Φ of Proposition 3, we obtain homomorphisms (of $\mathfrak{h}$-modules)

$$\omega(z_1, \dots, z_n, \lambda) : V^*_{\Lambda_1} \otimes \dots \otimes V^*_{\Lambda_n} \to F_{a_1,\dots,a_n}(z_1, \dots, z_n, \lambda)$$

which are isomorphisms for generic values of $z_1, \dots, z_n, \lambda$. The restriction of the map $\omega(z_1, \dots, z_n, \lambda)$ to $\mathbb{C}e^*_{m_1} \otimes \dots \otimes e^*_{m_n}$ is

$$\Phi_n(\omega(z_1, \lambda) \otimes \omega(z_2, \lambda - 2\eta\mu_1) \otimes \dots \otimes \omega(z_n, \lambda - 2\eta(\mu_1 + \dots + \mu_{n-1}))),$$

where $\mu_j = \Lambda_j - 2m_j$, $j = 1, \dots, n$. For example, if $n = 2$, then $\omega(z_1, z_2, \lambda)$ sends $e^*_j \otimes e^*_k$ to

$$\frac{1}{j!k!}\mathrm{Sym}\Big(\omega_j(t_1, \dots, t_j,\lambda; z_1)\omega_k(t_{j+1}, \dots, t_{j+k},\lambda - 2a_1 + 4\eta j; z_2) \prod_{i=j+1}^{j+k} \frac{\theta(t_i - z_1 + a_1)}{\theta(t_i - z_1 - a_1)}\Big),$$

where $\{\omega_j(t_1, \dots, t_j, \lambda; z)\}$ is the basis (4) of $F_a(z, \lambda)$.

More generally, we have an explicit formula for the image of $e^*_{m_1} \otimes \dots \otimes e^*_{m_n}$, which we discuss next.

3.3. **A basis of $F_{a_1,\dots,a_n}(z_1,\dots,z_n)$.** The space V_Λ comes with a basis e_j. Thus we have the natural basis $e^*_{m_1}\otimes\cdots\otimes e^*_{m_n}$ of the tensor product of $V^*_{\Lambda_i}$ in terms of the dual bases of the factors. The map $\omega(z_1,\dots,z_n,\lambda)$ maps, for generic z_i, this basis to a basis of $F_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$, which is an essential part of our formulae for Bethe ansatz eigenvectors and integral representations for solutions of the qKZB equations.

We give here an explicit formula for the basis vectors.

Proposition 4. *Let $m\in\mathbb{Z}_{\geqslant 0}$, $\Lambda=(\Lambda_1,\dots,\Lambda_n)\in\mathbb{C}^n$, and let $z=(z_1,\dots,z_n)\in\mathbb{C}^n$ be generic. Set $a_i=\eta\Lambda_i$. Let*

$$u(t_1,\dots,t_m)=\prod_{i<j}\frac{\theta(t_i-t_j+2\eta)}{\theta(t_i-t_j)}$$

Then, for generic $\lambda\in\mathbb{C}$, the functions

$$\omega_{m_1,\dots,m_n}(t_1,\dots,t_m,\lambda;z)=\omega(z,\lambda)\,e^*_{m_1}\otimes\cdots\otimes e^*_{m_n}$$

labeled by $m_1,\dots,m_n\in\mathbb{Z}$ with $\sum_k m_k=m$ form a basis of $F^m_a(z,\lambda)$ and are given by the explicit formula

$$\omega_{m_1,\dots,m_n}(t_1,\dots,t_m,\lambda;z)=u(t_1,\dots,t_m)^{-1}\sum_{I_1,\dots,I_n}\prod_{l=1}^{n}\prod_{i\in I_l}\prod_{k=1}^{l-1}\frac{\theta(t_i-z_k+a_k)}{\theta(t_i-z_k-a_k)}$$
$$\times\prod_{k<l}\prod_{i\in I_k,j\in I_l}\frac{\theta(t_i-t_j+2\eta)}{\theta(t_i-t_j)}\prod_{k=1}^{n}\prod_{j\in I_k}\frac{\theta(\lambda+t_j-z_k-a_k+2\eta m_k-2\eta\sum_{l=1}^{k-1}(\Lambda_l-2m_l))}{\theta(t_j-z_k-a_k)}.$$

The summation is over all n-tuples $I_1,\dots,I_n$ of disjoint subsets of $\{1,\dots,m\}$ such that I_j has m_j elements, $1\leqslant j\leqslant n$.

Proof: If $n=1$ this formula is the same as the one given above, eq. (4). The function u is designed to intertwine between the twisted symmetrization Sym defined above and the ordinary symmetrization $\mathrm{Sym}_0: f\mapsto\sum_{\sigma\in S_m}f(\sigma t)$. Indeed, for any function f of $t_1,\dots,t_m$, we have $\mathrm{Sym}_0(uf)=u\,\mathrm{Sym}(f)$. Let $I^0_1=\{1,\dots,m_1\}$, $I^0_2=\{m_1+1,\dots,m_1+m_2\}$ and so on. Also, if $I=\{i_1<\cdots<i_k\}$ set $t_I=(t_{i_1},\dots,t_{i_k})$. Then, by definition, $\omega_{m_1,\dots,m_n}$ is given by

$$\frac{1}{m_1!\cdots m_n!}\mathrm{Sym}\left(\prod_{k=1}^{n}\omega_{m_k}(t_{I^0_k},\lambda-2\eta\sum_{l=1}^{k-1}(\Lambda_l-2m_l);z_k)\prod_{l=1}^{k-1}\prod_{j\in I^0_l}\frac{\theta(t_j-z_l+a_l)}{\theta(t_j-z_l-a_l)}\right).$$

The one-point functions ω_{m_k} are given in eq. (4). Therefore, $(\prod m_j!)u\,\omega_{m_1,\dots,m_n}$ can be computed by ordinary symmetrization of the product of u with the expression in the brackets. It is then straightforward to compute this product, which is given by the term with $I_j=I^0_j$ in the sum above. The symmetrization gives all terms in the sum, each with multiplicity $\prod m_j!$. Thus the factorials are canceled and we get the claimed formula. □

3.4. ***R*-matrices.** Let $a=\eta\Lambda$ and $b=\eta\mathrm{M}$ be complex numbers. Since $F_{ab}(z,w,\lambda)=F_{ba}(w,z,\lambda)$ by the symmetry of the definition, we obtain a family of isomorphisms between $V^*_\Lambda\otimes V^*_\mathrm{M}$ and $V^*_\mathrm{M}\otimes V^*_\Lambda$. The composition of this family with the flip $P: V^*_\mathrm{M}\otimes V^*_\Lambda\to V^*_\Lambda\otimes V^*_\mathrm{M}$, $Pv\otimes w=w\otimes v$ gives a family of automorphisms of $V^*_\Lambda\otimes V^*_\mathrm{M}$:

Definition: Let z,w,λ be such that $\omega(z,w,\lambda): V^*_\Lambda\otimes V^*_\mathrm{M}\to F_{ab}(z,w,\lambda)$ is invertible. The *R-matrix* $R_{\Lambda,\mathrm{M}}(z,w,\lambda)\in\mathrm{End}_{\mathfrak{h}}(V_\Lambda\otimes V_\mathrm{M})$ is the dual map to the composition

$R^*_{\Lambda,\mathrm{M}}(z,w,\lambda)$:

$$V^*_\Lambda\otimes V^*_{\mathrm{M}}\xrightarrow{P}V^*_{\mathrm{M}}\otimes V^*_\Lambda\xrightarrow{\omega(w,z,\lambda)}F_{ab}(z,w,\lambda)\xrightarrow{\omega(z,w,\lambda)^{-1}}V^*_\Lambda\otimes V^*_{\mathrm{M}},$$

where we identify canonically $V^*_\Lambda\otimes V^*_{\mathrm{M}}$ with $(V_\Lambda\otimes V_{\mathrm{M}})^*$.

Alternatively, the R-matrix $R_{\Lambda,\mathrm{M}}(z,w,\lambda)$ can be thought of as a transition matrix expressing the basis $\tilde\omega_{ij}=\omega(w,z,\lambda)e^*_j\otimes e^*_i$ in terms of the basis $\omega_{ij}=\omega(z,w,\lambda)e^*_i\otimes e^*_j$ of $F_{ab}(z,w,\lambda)$: if $R_{\Lambda,\mathrm{M}}(z,w,\lambda)e_i\otimes e_j=\sum_{kl}R^{kl}_{ij}e_k\otimes e_l$, then

$$\tilde\omega_{kl}=\sum_{ij}R^{kl}_{ij}\omega_{ij}.$$

Remark. Note that by the symmetry under reversal of all signs,

$$R_{\Lambda,\mathrm{M}}(z,\lambda)=R_{-\Lambda,-\mathrm{M}}(-z,-\lambda),$$

if we identify V_Λ with $V_{-\Lambda}$ via the basis (e_j).

Lemma 5. (i) *$R_{\Lambda,\mathrm{M}}(z,w,\lambda)$ is a meromorphic function of $\Lambda,\mathrm{M},z,w,\lambda$.*

(ii) *If Λ is generic, then $R_{\Lambda,\Lambda}(z,w,\lambda)$ is regular at $z=w$ and $\lim_{z\to w}R_{\Lambda,\Lambda}(z,w,\lambda)=P$, the flip $u\otimes v\mapsto v\otimes u$.*

(iii) *$R_{\Lambda,\mathrm{M}}(z,w,\lambda)$ depends only on the difference $z-w$.*

Proof: (i) Consider the bases $\{\omega_{ij}\}$, $\{\tilde\omega_{ij}\}$, $i+j=m$, of $F^m_{ab}(z,w,\lambda)$. These functions are meromorphic in all variables. Since they are linearly independent there are $m+1$ values $t^0,\dots,t^m$ of $t\in\mathbb{C}^m$ so that $(\omega_{i,m-i}(t^j))_{0\leqslant i,j\leqslant m}$ is an invertible matrix B. The R-matrix is then $B^{-1}\tilde B$ where $\tilde B$ is the matrix with entries $\tilde\omega_{i,m-i}(t^j)$. Therefore it is meromorphic.
(ii) The proof of this is deferred to the end of Section 8.
(iii) We have, for all $c\in\mathbb{C}$, an isomorphism $U_c: F_{ab}(z,w,\lambda)\to F_{ab}(z+c,w+c,\lambda)$, given by the translation of the arguments t_i by c. It is easy to see that $\omega(z+c,w+c,\lambda)U_c=\omega(z,w,\lambda)$, so $R_{\Lambda,\mathrm{M}}(z+c,w+c,\lambda)=R_{\Lambda,\mathrm{M}}(z,w,\lambda)$. □

Accordingly, we write $R_{\Lambda,\mathrm{M}}(z-w,\lambda)$ instead of $R_{\Lambda,\mathrm{M}}(z,w,\lambda)$ in what follows.

By definition the R-matrix is determined by the relation $\omega(z_1,z_2,\lambda)R^*_{\Lambda,\mathrm{M}}(z_1-z_2,\lambda)=\omega(z_2,z_1,\lambda)P$ in $\mathrm{Hom}(V^*_\Lambda\otimes V^*_{\mathrm{M}},F_{ab}(z_1,z_2,\lambda))$. More generally, by the associativity of Φ, we have:

Lemma 6. *For any $\Lambda_1,\dots,\Lambda_n$, the identity*

$$\begin{aligned}\omega(\dots,z_j,z_{j+1},\dots,\lambda)\,R^*_{\Lambda_j\Lambda_{j+1}}(z_j-z_{j+1},\lambda-2\eta\textstyle\sum_{l<j}h^{(l)})^{(j,j+1)}\\ =\omega(\dots,z_{j+1},z_j,\dots,\lambda)P^{(j,j+1)}\end{aligned}$$

holds in $\mathrm{Hom}(V^*_{\Lambda_1}\otimes\cdots\otimes V^*_{\Lambda_n},F_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda))$, $a_i=\eta\Lambda_i$.

A corollary of this lemma is the dynamical Yang–Baxter equation for the R-matrices:

Theorem 7. *The matrices $R_{\Lambda,\mathrm{M}}(z,\lambda)$ obey I–III of Section 2.*

Proof: Fix z_i,Λ_i, $i=1,2,3$ and let $R_{ij}(\lambda)=R_{\Lambda_i,\Lambda_j}(z_i-z_j,\lambda)$. The dynamical Yang–Baxter equation is equivalent to the equation

$$R^{*(23)}_{23}(\lambda-2\eta h^{(1)})R^{*(13)}_{13}(\lambda)R^{*(12)}_{12}(\lambda-2\eta h^{(3)})=R^{*(12)}_{12}(\lambda)R^{*(13)}_{13}(\lambda-2\eta h^{(2)})R^{*(23)}_{23}(\lambda)$$

for the dual maps $R^*_{ij}(\lambda)$. The latter equation follows from using the previous lemma several times to express $\omega(z_3,z_2,z_1,\lambda)$ in terms of $\omega(z_1,z_2,z_3,\lambda)$ in two different ways.

The second claim follows by expressing $\omega(w,z,\lambda)$ in terms of $\omega(z,w,\lambda)$ and conversely. □

Let us now consider the case of positive integer weights. In this case the R-matrices have invariant subspaces. If $\Lambda\in\mathbb{Z}_{\geqslant 0}$ we let SV_Λ be the subspace of V_Λ spanned by $e_{\Lambda+1}$, $e_{\Lambda+2},\dots$. The $\Lambda+1$-dimensional quotient V_Λ/SV_Λ will be denoted by L_Λ, and will be often identified with $\oplus_{j=0}^m\mathbb{C}e_j$.

Theorem 8. *Let z,η,λ be generic and Λ, $\mathrm{M}\in\mathbb{C}$.*

(i) *If $\Lambda\in\mathbb{Z}_{\geqslant 0}$, then $R_{\Lambda,\mathrm{M}}(z,\lambda)$ preserves $SV_\Lambda\otimes V_\mathrm{M}$*
(ii) *If $\mathrm{M}\in\mathbb{Z}_{\geqslant 0}$, then $R_{\Lambda,\mathrm{M}}(z,\lambda)$ preserves $V_\Lambda\otimes SV_\mathrm{M}$*
(iii) *If $\Lambda\in\mathbb{Z}_{\geqslant 0}$ and $\mathrm{M}\in\mathbb{Z}_{\geqslant 0}$, then $R_{\Lambda,\mathrm{M}}(z,\lambda)$ preserves $SV_\Lambda\otimes V_\mathrm{M}+V_\Lambda\otimes SV_\mathrm{M}$.*

Proof: The claims of this theorem are equivalent to the statements that the operators dual to the R-matrices preserve the subspaces $L_\Lambda^*\otimes V_\mathrm{M}^*$, $V_\Lambda^*\otimes L_\mathrm{M}^*$, $L_\Lambda^*\otimes L_\mathrm{M}^*$, respectively. This follows from Theorem 31 below, which gives a characterization of the images by $\omega(z,w,\lambda)$ of these subspaces in terms of residue conditions. □

In particular, if Λ and/or M are nonnegative integers, $R_{\Lambda,\mathrm{M}}(z,\lambda)$ induces operators, still denoted by $R_{\Lambda,\mathrm{M}}(z,\lambda)$, on the quotients $L_\Lambda\otimes V_\mathrm{M}$, $V_\Lambda\otimes L_\mathrm{M}$ and/or $L_\Lambda\otimes L_\mathrm{M}$. They obey the dynamical Yang–Baxter equation.

3.5. **Examples.** We give now some examples of computations of matrix elements of the R-matrix $R_{\Lambda,\mathrm{M}}(z-w,\lambda)$, assuming that the parameters are generic.

The R-matrix is calculated as the transition matrix relating two bases of $F_{ab}(z,w,\lambda)$: let

$$\tilde\omega_{ij}=\omega(w,z,\lambda)\,e_j^*\otimes e_i^*,\qquad \omega_{ij}=\omega(z,w,\lambda)\,e_i^*\otimes e_j^*.$$

The matrix elements of R with respect to the basis $e_j\otimes e_k$ are given by $\tilde\omega_{kl}=\sum_{ij}R_{ij}^{kl}\omega_{ij}$. By construction, this matrix commutes with $h^{(1)}+h^{(2)}$, and thus preserves the weight spaces

$$(V_\Lambda\otimes V_\mathrm{M})[\Lambda+\mathrm{M}-2m]=\oplus_{j=0}^m\mathbb{C}e_j\otimes e_{m-j}$$

We may therefore consider the problem of computing the matrix elements of the R-matrix separately on each weight space. Without loss of generality we assume that $w=0$.

Let $m=0$. Then the weight space is spanned by $e_0\otimes e_0$ and $\omega_{00}=\tilde\omega_{00}=1$. Therefore $R_{00}^{00}=1$.

Let now $m=1$. The basis elements are functions of one variable $t=t_1$ and we have (with $a=\eta\Lambda$, $b=\eta\mathrm{M}$)

$$\omega_{01}(t)=\frac{\theta(\lambda+2\eta+t-2a-b)\theta(t-z+a)}{\theta(t-b)\theta(t-z-a)},\qquad \omega_{10}(t)=\frac{\theta(\lambda+2\eta+t-z-a)}{\theta(t-z-a)},$$

and

$$\tilde\omega_{01}(t)=\frac{\theta(\lambda+2\eta+t-b)}{\theta(t-b)},\qquad \tilde\omega_{10}(t)=\frac{\theta(\lambda+2\eta+t-z-2b-a)\theta(t+b)}{\theta(t-z-a)\theta(t-b)}.$$

To express one basis in terms of the other we notice that these functions have simple poles at two points (modulo the lattice). The elements of the R-matrix are determined by comparing the residues at these two points. We have

$$\operatorname{res}_{t=b}\omega_{01}(t)=\frac{\theta(\lambda+2\eta-2a)\theta(z-b-a)}{\theta'(0)\theta(z-b+a)},\qquad \operatorname{res}_{t=z+a}\omega_{01}(t)=\frac{\theta(\lambda+2\eta+z-a-b)\theta(2a)}{\theta'(0)\theta(z+a-b)},$$

$$\operatorname{res}_{t=b}\omega_{10}(t)=0,\qquad \operatorname{res}_{t=z+a}\omega_{10}(t)=\frac{\theta(\lambda+2\eta)}{\theta'(0)}.$$

On the other hand,

$$\operatorname{res}_{t=b}\tilde\omega_{01}(t)=\frac{\theta(\lambda+2\eta)}{\theta'(0)},\qquad \operatorname{res}_{t=z+a}\tilde\omega_{01}(t)=0,$$

$$\operatorname{res}_{t=b}\tilde\omega_{10}(t)=\frac{\theta(\lambda+2\eta-z-b-a)\theta(2b)}{\theta'(0)\theta(-z+b-a)},\qquad \operatorname{res}_{t=z+a}\tilde\omega_{10}(t)=\frac{\theta(\lambda+2\eta-2b)\theta(z+a+b)}{\theta'(0)\theta(z+a-b)}.$$

If we put these residues into matrices $\begin{pmatrix}A & B\\ 0 & D\end{pmatrix}$ and $\begin{pmatrix}\tilde A & 0\\ \tilde C & \tilde D\end{pmatrix}$, respectively, we get that the restriction of $R_{\Lambda,\mathrm{M}}(z,\lambda)$ to $(V_\Lambda\otimes V_{\mathrm{M}})[\Lambda+\mathrm{M}-2]$ in the basis $e_0\otimes e_1$, $e_1\otimes e_0$ is given by the matrix

$$\begin{pmatrix}\tilde A & 0\\ \tilde C & \tilde D\end{pmatrix}\begin{pmatrix}A & B\\ 0 & D\end{pmatrix}^{-1}=\begin{pmatrix}\tilde A A^{-1} & -\tilde A B A^{-1}D^{-1}\\ \tilde C A^{-1} & -\tilde C B A^{-1} D^{-1}+\tilde D D^{-1}\end{pmatrix}.$$

These matrix elements have product form except the last one, which can be further simplified:

$$\begin{aligned}R^{10}_{10} &= -\tilde C B A^{-1}D^{-1}+\tilde D D^{-1}\\ &= \frac{\theta(\lambda+2\eta-z-a-b)\theta(2b)\theta(\lambda+2\eta+z-a-b)\theta(2a)}{\theta(\lambda+2\eta)\theta(z-a-b)\theta(\lambda+2\eta-2a)\theta(z+a-b)}\\ &\quad+\frac{\theta(\lambda+2\eta-2b)\theta(z+a+b)}{\theta(\lambda+2\eta)\theta(z+a-b)}.\end{aligned}$$

To simplify this expression, consider it as a function of λ: it is a 1-periodic function, which gets multiplied by $e^{4\pi ib}$ when λ is replaced by $\lambda+\tau$. It is regular at the apparent singularity $\lambda=-2\eta$. So it has only simple poles at $\lambda=2\eta-2a$ and its translates by $\mathbb{Z}+\tau\mathbb{Z}$. These properties uniquely define this function up to a multiplicative function: it can therefore be written in the form $C\theta(\lambda+2\eta-2a-2b)/\theta(\lambda+2\eta-2a)$. The constant C can be determined by computing the residue at $\lambda=2a-2\eta$.

Let us summarize the results of these calculations.

Proposition 9. *Let the matrix elements of the R-matrix $R_{\Lambda,\mathrm{M}}(z,\lambda)$ be defined by*

$$R_{\Lambda,\mathrm{M}}(z,\lambda)e_i\otimes e_j=\textstyle\sum_{kl}R^{kl}_{ij}e_k\otimes e_l.$$

Then

$$\begin{aligned}R^{00}_{00} &= 1\\ R^{01}_{01} &= \frac{\theta(z+\eta\Lambda-\eta\mathrm{M})\theta(\lambda+2\eta)}{\theta(z-\eta\Lambda-\eta\mathrm{M})\theta(\lambda+2\eta(1-\Lambda))}\\ R^{01}_{10} &= -\frac{\theta(\lambda+2\eta+z-\eta\Lambda-\eta\mathrm{M})\theta(2\eta\Lambda)}{\theta(z-\eta\Lambda-\eta\mathrm{M})\theta(\lambda+2\eta(1-\Lambda))}\\ R^{10}_{01} &= -\frac{\theta(\lambda+2\eta-z-\eta\Lambda-\eta\mathrm{M})\theta(2\eta\mathrm{M})}{\theta(z-\eta\Lambda-\eta\mathrm{M})\theta(\lambda+2\eta(1-\Lambda))},\\ R^{10}_{10} &= \frac{\theta(z+\eta\mathrm{M}-\eta\Lambda)\theta(\lambda+2\eta(1-\Lambda-\mathrm{M}))}{\theta(z-\eta\Lambda-\eta\mathrm{M})\theta(\lambda+2\eta(1-\Lambda))}.\end{aligned}$$

For higher values of m, the matrix elements can still be computed along the same lines. The residues are replaced by suitable iterated residues defined for functions of several variables. This technique is explained in Section 8. The iterated residues of basis elements ω_{ij} and $\tilde{\omega}_{ij}$ form triangular matrices whose diagonal matrix elements do not vanish for generic parameters. This gives an expression of the restriction of the R-matrix to any weight space as a product of a lower triangular matrix by an upper triangular matrix, whose entries are iterated residues, just as in the one-variable case.

In the next subsection we present the results of this calculation relevant for the representation theory of elliptic quantum groups.

3.6. **Evaluation Verma modules and their tensor products.** Here we explain the relation between the geometric construction of tensor products and R-matrices and the representation theory of $E_{\tau,\eta}(sl_2)$ [FV1].

We first recall the definition of a representation of $E_{\tau,\eta}(sl_2)$: let $\mathfrak{h}$ act on $\mathbb{C}^2$ via $h = \mathrm{diag}(1,-1)$. A representation of $E_{\tau,\eta}(sl_2)$ is an $\mathfrak{h}$-module W with diagonalizable action of h and finite dimensional eigenspaces, together with an operator $L(z,\lambda) \in \mathrm{End}(\mathbb{C}^2 \otimes W)$ (the "L-operator"), commuting with $h^{(1)} + h^{(2)}$, and obeying the relations

$$\begin{aligned} R^{(12)}(z-w,\lambda-2\eta h^{(3)})\ & L^{(13)}(z,\lambda)\, L^{(23)}(w,\lambda-2\eta h^{(1)}) \\ &= L^{(23)}(w,\lambda)\, L^{(13)}(z,\lambda-2\eta h^{(2)}) R^{(12)}(z-w,\lambda) \end{aligned}$$

in $\mathrm{End}(\mathbb{C}^2\otimes\mathbb{C}^2\otimes W)$. The *fundamental R-matrix* $R(z,\lambda) \in \mathrm{End}(\mathbb{C}^2\otimes\mathbb{C}^2)$ is the following solution of the dynamical Yang–Baxter equation: let e_0, e_1 be the standard basis of $\mathbb{C}^2$, then with respect to the basis $e_0\otimes e_0$, $e_0\otimes e_1$, $e_1\otimes e_0$, $e_1\otimes e_1$ of $\mathbb{C}^2\otimes\mathbb{C}^2$,

$$R(z,\lambda) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \alpha(z,\lambda) & \beta(z,\lambda) & 0 \\ 0 & \beta(z,-\lambda) & \alpha(z,-\lambda) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

where

$$\alpha(z,\lambda) = \frac{\theta(\lambda+2\eta)\theta(z)}{\theta(\lambda)\theta(z-2\eta)}, \qquad \beta(z,\lambda) = -\frac{\theta(\lambda+z)\theta(2\eta)}{\theta(\lambda)\theta(z-2\eta)}.$$

To discuss representation theory, it is convenient to think of $L(z,\lambda) \in \mathrm{End}(\mathbb{C}^2\otimes V)$ as a two by two matrix with entries $a(z,\lambda)$, $b(z,\lambda)$, $c(z,\lambda)$, $d(z,\lambda)$ in $\mathrm{End}(W)$. In [FV1] we wrote explicitly the relations that these four operators must satisfy, and defined a class of representations, the evaluation Verma modules, by giving explicitly the action of these four operators on basis vectors. Next, we show how these formulae can be obtained from our geometric construction.

Theorem 10. *Let us identify L_1 with $\mathbb{C}^2$ via the basis e_0, e_1. The R-matrix $R_{1,1}(z,\lambda) \in \mathrm{End}(L_1\otimes L_1)$ coincides with the fundamental R-matrix.*

Proof: Most of the matrix elements have been computed in 3.5. The only remaining element to compute is R^{11}_{11}. By Theorem 8, we know that the dual of the R-matrix preserves the one-dimensional space spanned by $e_1^*\otimes e_1^*$ if $\Lambda = \mathrm{M} = 1$. This means that $\omega_{1,1} = \omega(z,w,\lambda)e_1^*\otimes e_1^*$ is proportional to $\tilde{\omega}_{1,1} = \omega(w,z,\lambda)e_1^*\otimes e_1^*$. We have, by definition,

$$\begin{aligned} \omega_{1,1} \;=\;& \frac{\theta(\lambda+t_1+\eta-z)\theta(\lambda+t_2+3\eta-w)\theta(t_2-z+\eta)}{\theta(t_1-z-\eta)\theta(t_2-w-\eta)\theta(t_2-z-\eta)} \\ &+\frac{\theta(\lambda+t_2+\eta-z)\theta(\lambda+t_1+3\eta-w)\theta(t_1-z+\eta)\theta(t_1-t_2-2\eta)}{\theta(t_2-z-\eta)\theta(t_1-w-\eta)\theta(t_1-z-\eta)\theta(t_1-t_2+2\eta)}. \end{aligned}$$

The formula for $\tilde{\omega}_{1,1}$ is obtained by interchanging z and w. To find the constant of proportionality (the inverse of R_{11}^{11}), we compute both expressions at some point. We can take for instance $t_1 = 2\eta$, $t_2 = 0$. Then it is obvious that at this point $\omega_{1,1} = \tilde{\omega}_{1,1}$. Thus $R_{11}^{11} = 1$. $\square$

Corollary 11. *For any $w, \mathrm{M} \in \mathbb{C}$, the $\mathfrak{h}$-module V_{M} together with the operator $L(z,\lambda) = R_{1,\mathrm{M}}(z-w,\lambda) \in \mathrm{End}(L_1 \otimes V_{\mathrm{M}})$ defines a representation of $E_{\tau,\eta}(sl_2)$.*

This representation is called in [FV1] the evaluation Verma module with evaluation point w and highest weight M. It is denoted by $V_{\mathrm{M}}(w)$. The matrix elements of $L(z,\lambda)$ can be computed straightforwardly by the method described in the previous section. The result is given explicitly in [FV1], Theorem 3, in terms of the action of $a(z,\lambda), \dots, d(z,\lambda)$. In the notation of Proposition 9, this result amounts to the following formulae for the matrix elements R_{ij}^{kl} of $R_{1,\Lambda}(z,\lambda) \in \mathrm{End}(L_1 \otimes V_\Lambda)$.

$$
\begin{aligned}
R_{0k}^{0k} &= \frac{\theta(z-(\Lambda+1-2k)\eta)}{\theta(z-(\Lambda+1)\eta)}\frac{\theta(\lambda+2k\eta)}{\theta(\lambda)},\\
R_{1k}^{0,k+1} &= -\frac{\theta(\lambda+z-(\Lambda-1-2k)\eta)}{\theta(z-(\Lambda+1)\eta)}\frac{\theta(2\eta)}{\theta(\lambda)},\\
R_{0k}^{1,k-1} &= -\frac{\theta(\lambda-z-(\Lambda+1-2k)\eta)}{\theta(z-(\Lambda+1)\eta)}\frac{\theta(2(\Lambda+1-k)\eta)}{\theta(\lambda)}\frac{\theta(2k\eta)}{\theta(2\eta)},\\
R_{1k}^{1k} &= \frac{\theta(z-(-\Lambda+1+2k)\eta)}{\theta(z-(\Lambda+1)\eta)}\frac{\theta(\lambda-2(\Lambda-k)\eta)}{\theta(\lambda)}.
\end{aligned}
$$

These formulae can be obtained by computing residues, as explained in the previous subsection. Moreover, the tensor product construction of 3.2 is related to the tensor product of representations of the elliptic quantum group. Recall that if W_1, W_2 are representations of the elliptic quantum group with L-operators $L_1(z,\lambda)$, $L_2(z,\lambda)$, then their tensor product $W = W_1 \otimes W_2$ with L-operator

$$L(z,\lambda) = L_1(z,\lambda-2\eta h^{(3)})^{(12)} L_2(z,\lambda)^{(13)} \in \mathrm{End}(\mathbb{C}^2 \otimes W)$$

is also a representation of the elliptic quantum group.

Theorem 12. *Let $\Lambda_1, \dots, \Lambda_n \in \mathbb{C}$ and $z_1, \dots, z_n$ be generic complex numbers. Let $V = V_{\Lambda_1} \otimes \cdots \otimes V_{\Lambda_n}$ and $L(z,\lambda) \in \mathrm{End}(V_{\Lambda=1} \otimes V)$ be defined by the relation*

$$\omega(z, z_1, \dots, z_n, \lambda) L(z,\lambda)^* = \omega(z_1, \dots, z_n, z, \lambda) P$$

in $\mathrm{End}((V_1 \otimes V)^*) = \mathrm{End}(V_1^* \otimes V^*)$, where $P v_1 \otimes v = v \otimes v_1$, if $v_1 \in V_1^*$, $v \in V^*$. Then $L(z,\lambda)$ is well-defined as an endomorphism of the quotient $L_1 \otimes V = \mathbb{C}^2 \otimes V$, and defines the structure of a representation of $E_{\tau,\eta}(sl_2)$ on V. This representation is isomorphic to the tensor product of evaluation Verma modules

$$V_{\Lambda_n}(z_n) \otimes \cdots \otimes V_{\Lambda_1}(z_1),$$

with isomorphism $u_1 \otimes \cdots \otimes u_n \mapsto u_n \otimes \cdots \otimes u_1$

Proof: We prove this if $n = 2$. The proof in the general case is the same, but requires more writing.

By applying Lemma 6 twice, we get

$$
\begin{aligned}
&\omega(z, z_1, z_2, \lambda) R_{1,\Lambda_1}^*(z-z_1,\lambda)^{(12)} R_{1,\Lambda_2}^*(z-z_2,\lambda-2\eta h^{(2)})^{(13)}\\
&= \omega(z_1, z, z_2, \lambda) R_{1,\Lambda_2}^*(z-z_2,\lambda-2\eta h^{(1)})^{(23)} P^{(12)}\\
&= \omega(z_1, z_2, z, \lambda) P^{(23)} P^{(12)}.
\end{aligned}
$$

But $P^{(23)}P^{(12)} = P$, the flip $V_1 \otimes V \to V \otimes V_1$. By taking duals, we obtain

$$L(z,\lambda) = R_{1,\Lambda_2}(z - z_2, \lambda - 2\eta h^{(2)})^{(13)} R_{1,\Lambda_1}(z - z_1, \lambda)^{(12)}.$$

This operator in well-defined as an endomorphism of the quotient $L_1 \otimes V$, since its factors are. Moreover

$$P^{(23)} L(z,\lambda) P^{(23)} = R_{1,\Lambda_2}(z - z_2, \lambda - 2\eta h^{(3)})^{(12)} R_{1,\Lambda_1}(z - z_1, \lambda)^{(13)},$$

which is the L-operator of the representation $V_{\Lambda_2}(z_2) \otimes V_{\Lambda_1}(z_1)$. □

Finally, the dynamical Yang–Baxter equation in $L_1 \otimes V_\Lambda \otimes V_{\mathrm{M}}$ can be stated as saying that $R_{\Lambda,\mathrm{M}}(z-w,\lambda)P$ is an isomorphism from $V_{\mathrm{M}}(w) \otimes V_\Lambda(z)$ to $V_\Lambda(z) \otimes V_{\mathrm{M}}(w)$, see [FV1]. Therefore, by uniqueness, the R-matrices constructed here coincide with the solutions of the dynamical Yang–Baxter equation described in Section 13 of [FV1].

4. Bethe ansatz

In this section we fix the parameters η, τ of the elliptic quantum group and n complex numbers $\Lambda_1, \dots, \Lambda_n$ such that $\sum \Lambda_l = 2m$ for some nonnegative integer m. We also set $a_j = \eta \Lambda_j$, $j = 1, \dots, n$. Let $z_1, \dots, z_n$ be generic complex numbers. Then the zero weight space $V[0]$ of $V = V_{\Lambda_1} \otimes \cdots \otimes V_{\Lambda_n}$ is non-trivial, and we have commuting difference operators $H_j = H_j(z_1, \dots, z_n)$, $j = 1, \dots, n$, cf. (2), acting on the space $\mathrm{Fun}(V[0])$.

We give a formula for common quasiperiodic eigenfunctions of the operators H_j, i.e., functions ψ such that $H_j \psi = \epsilon_j \psi$, $j = 1, \dots, n$, for some $\epsilon_j \in \mathbb{C}$. The quasiperiodicity assumption means that we require that $\psi(\lambda + 1) = \mu \psi(\lambda)$ for some multiplier $\mu \in \mathbb{C}^\times$.

Our formula is given in terms of the expressions of the previous section. For $j_1, \dots, j_n \in \mathbb{Z}_{\geqslant 0}$ such that $\sum j_i = m$, let $\omega_{j_1,\dots,j_n}(t_1, \dots, t_m, \lambda)$ be the value at $t_1, \dots, t_m$ of the function $\omega(z_1, \dots, z_n, \lambda) e^*_{j_1} \otimes \cdots \otimes e^*_{j_n}$.

Theorem 13. *(cf.* [TV1]*) Let $c \in \mathbb{C}$. Suppose that $t_1, \dots, t_m$ obey the system of "Bethe ansatz" equations*

$$\prod_{l=1}^{n} \frac{\theta(t_j - z_l + a_l)}{\theta(t_j - z_l - a_l)} \prod_{k: k \neq j} \frac{\theta(t_j - t_k - 2\eta)}{\theta(t_j - t_k + 2\eta)} = e^{-4\eta c}, \qquad j = 1, \dots, m. \tag{5}$$

Then

$$\psi(\lambda) = \sum_{j_1 + \cdots + j_n = m} e^{c\lambda} \omega_{j_1,\dots,j_n}(t_1, \dots, t_m, \lambda)\, e_{j_1} \otimes \cdots \otimes e_{j_n} \tag{6}$$

is a common eigenfunction of the commuting operators H_j, $j = 1, \dots, n$, with eigenvalues

$$\epsilon_j = e^{-2ca_j} \prod_{k=1}^{m} \frac{\theta(t_k - z_j - a_j)}{\theta(t_k - z_j + a_j)}$$

and multiplier $\mu = (-1)^m e^c$. Moreover, if $t_1, \dots, t_n$ are a solution of (5), and $\sigma \in S_m$ is any permutation, then $t_{\sigma(1)}, \dots, t_{\sigma(n)}$ are also a solution. The eigenfunctions ψ corresponding to these two solutions are proportional.

Example: Let $m = 1$. Thus $\Lambda_1 + \cdots + \Lambda_n = 2$. The Bethe ansatz equation for $t = t_1$ and c is

$$\prod_{l=1}^{n} \frac{\theta(t - z_l - \eta\Lambda_l)}{\theta(t - z_l + \eta\Lambda_l)} = e^{4\eta c}.$$

If t, c obey this equation, then

$$\psi(\lambda) = e^{c\lambda} \sum_{l=1}^{n} \left(\frac{\theta(\lambda + t - z_l - \eta\Lambda_l - 2\eta \sum_{j=1}^{l-1} \Lambda_j)}{\theta(t - z_l - \eta\Lambda_l)} \prod_{j=1}^{l-1} \frac{\theta(t - z_j + \eta\Lambda_l)}{\theta(t - z_j - \eta\Lambda_l)} \right) e_0 \otimes \cdots \otimes e_1 \otimes \cdots \otimes e_0,$$

(e_1 is at the lth position) is a common eigenfunction of the difference operators $H_j(z)$ $j = 1, \ldots, n$. The eigenvalues are

$$\epsilon_j = e^{-2\eta c \Lambda_j} \frac{\theta(t - z_j - \eta\Lambda_j)}{\theta(t - z_j + \eta\Lambda_j)}.$$

The proof of this theorem occupies the rest of this section. The fact that solutions are mapped to solutions by permutations is clear from the form of the equations (5). The functions $\omega_{j_1,\ldots,j_n}$ are symmetric under the action of permutations of Lemma 2. Therefore functions $\psi(\lambda)$ corresponding to solutions related by permutations are proportional.

We are left to prove that ψ is an eigenfunction of the operators H_j.

We first reduce the problem to showing that $\psi(\lambda)$ is an eigenfunction of H_1 for all permutations of the parameters z_i, Λ_i. For this we must study the dependence of H_j on the ordering of the parameters z_i, Λ_i. Let us introduce some notation to indicate explicitly the dependence on the parameters: let S_m act on $\mathbb{C}^m$ by permutations of the coordinates, and let s_j be the transposition of the jth and $(j+1)$st coordinates. Let us write $H_j(\Lambda, z)$ for the operator H_j acting on the space of functions $\mathcal{H}(\Lambda)$ of functions of λ with values in $(V_{\Lambda_1} \otimes \cdots \otimes V_{\Lambda_n})[0]$, $\Lambda, z \in \mathbb{C}^n$.

Lemma 14. *Let $\Lambda = (\Lambda_1, \ldots, \Lambda_n)$, $z = (z_1, \ldots, z_n) \in \mathbb{C}^n$ and let $S_j : \mathcal{H}(\Lambda) \to \mathcal{H}(s_j\Lambda)$ $(j = 1, \ldots n-1)$ be the linear map*

$$S_j f(\lambda) = P^{(j,j+1)} R^{(j,j+1)}_{\Lambda_j \Lambda_{j+1}}(z_j - z_{j+1}, \lambda - 2\eta(h^{(1)} + \cdots + h^{(j-1)})) f(\lambda).$$

Then $H_{j+1}(s_j\Lambda, s_j z) S_j = S_j H_j(\Lambda, z)$, for all $j = 1, \ldots, n-1$.

Proof: The proof is a matter of writing the claim using the representation of H_j given in (2), and bringing $P^{(j,j+1)}$ to the left of the R-matrices, using commutation relations. □

Notice that the equations (5) do not depend on the ordering of the parameters z_i, Λ_i. Let us fix a solution t^* of (5) with parameters $\Lambda = (\Lambda_1, \ldots, \Lambda_n)$ and $z = (z_1, \ldots, z_n)$. Then for each permutation $\sigma \in S_n$, t^* is still a solution of the equations (5) with parameters $\sigma\Lambda$, σz and we have a corresponding function $\psi_\sigma(\lambda) \in \mathcal{H}(\sigma\Lambda)$ given by the formula (6) with parameters $\sigma\Lambda$, σz. It takes values in $V_{\Lambda_{\sigma^{-1}(1)}} \otimes \cdots \otimes V_{\Lambda_{\sigma^{-1}(n)}}$. We may now rephrase Lemma 6 as follows:

$$\psi_{s_j}(\lambda) = P^{(j,j+1)} R^{(j,j+1)}_{\Lambda_j \Lambda_{j+1}}(z_j - z_{j+1}, \lambda - 2\eta(h^{(1)} + \cdots + h^{(j-1)})) \psi(\lambda).$$

With Lemma 14, we see that the fact that $\psi(\lambda)$ is an eigenfunction of $H_j(\Lambda, z)$ with eigenvalue $\epsilon_j = \epsilon_j(\Lambda, z)$ for all $j = 1, \ldots, n$ is equivalent to the fact that $\psi_\sigma(\lambda)$ is an eigenfunction of $H_1(\sigma\Lambda, \sigma z)$ with eigenvalue $\epsilon_1(\sigma\Lambda, \sigma z)$ for all $\sigma \in S_n$.

Therefore, it is sufficient to prove that ψ is an eigenfunction of H_1 with eigenvalue ϵ_1 for any solution of (5) with arbitrary parameters.

Lemma 15. *Let $u \in V^*_{\Lambda_1}[\mu]$, $v \in (V^*_{\Lambda_2} \otimes \cdots \otimes V^*_{\Lambda_n})[-\mu]$. Thus $\mu = \Lambda_1 - 2m' = -\sum_{j=2}^{n} \Lambda_j + 2m''$ for some integers $m', m'' \geqslant 0$. Set $m = m' + m''$. Let $f = \omega(z_1, \lambda)u$,*

$g=\omega(z_2,\dots,z_n,\lambda-2\eta\mu)v$. Then $\omega(z_1,\dots,z_n,\lambda)$ maps $u\otimes v$ to

$$\frac{1}{(n-1)!}\mathrm{Sym}\Big(f(t_1,\dots,t_{m'})g(t_{m'+1},\dots,t_m)\prod_{j>m'}\frac{\theta(t_j-z_1+a_1)}{\theta(t_j-z_1-a_1)}\Big),$$

and $\omega(z_2,\dots,z_n,z_1,\lambda-2\eta\mu)$ maps $v\otimes u$ to

$$\frac{1}{(n-1)!}\mathrm{Sym}\Big(g(t_{m'+1},\dots,t_m)f(t_1,\dots,t_{m'})\prod_{\substack{j\leqslant m'\\ l\geqslant 2}}\frac{\theta(t_j-z_l+a_l)}{\theta(t_j-z_l-a_l)}\prod_{\substack{j\leqslant m'\\ k>m'}}\frac{\theta(t_j-t_k-2\eta)}{\theta(t_j-t_k+2\eta)}\Big).$$

Proof: The first formula is a rewriting of the definition. In the second formula, the expression appearing in the definition was replaced by another term in the sum over S_m defining Sym, which gives the same result. □

Lemma 16. *If $t^*=(t_1^*,\dots,t_m^*)$ is a solution of (5) and u, v are as in Lemma 15, then*

$$\omega(z_2,\dots,z_n,z_1,\lambda-2\eta\mu)v\otimes u|_{t^*}=e^{2c\eta\mu}\epsilon_1\cdot\omega(z_1,\dots,z_n,\lambda)u\otimes v|_{t^*}$$

Here, $|_{t^*}$ denotes the value at t^* of a function of $t_1,\dots,t_m$.

Proof: Let us introduce the ratio $h(t_1,\dots,t_m)$ of the factors appearing in the previous lemma:

$$\prod_{\substack{j\leqslant m'\\ l>2}}\frac{\theta(t_j-z_l+a_l)}{\theta(t_j-z_l-a_l)}\prod_{\substack{j\leqslant m'\\ k>m'}}\frac{\theta(t_j-t_k-2\eta)}{\theta(t_j-t_k+2\eta)}=h(t_1,\dots,t_m)\prod_{j>m'}\frac{\theta(t_j-z_1+a_1)}{\theta(t_j-z_1-a_1)}.$$

This equation can be rewritten as

$$h(t_1,\dots,t_m)=e^{2ca_1}\epsilon_1(t_1,\dots,t_m)\prod_{j\leqslant m'}\Big(\prod_{l=1}^{n}\frac{\theta(t_j-z_l-a_l)}{\theta(t_j-z_l+a_l)}\prod_{k>m'}\frac{\theta(t_j-t_k+2\eta)}{\theta(t_j-t_k-2\eta)}\Big),$$

where $\epsilon_1(t_1\dots,t_m)=e^{-2ca_1}\prod_{j=1}^m\theta(t_j-z_1-a_1)/\theta(t_j-z_1+a_1)$. The product of the first m' Bethe ansatz equations (5), implies that, for $t=t^*$,

$$h(t_1,\dots,t_m)=e^{2ca_1}e^{-4\eta cm'}\epsilon_1(t_1,\dots,t_m)$$

The right-hand side of this equation is symmetric under permutations of $t_1,\dots,t_m$. Moreover, if $t_1,\dots,t_m$ is a solution of (5) then also $t_{\sigma(1)},\dots,t_{\sigma(m)}$ for any permutation σ. We conclude that for all solutions of (5),

$$h(t_1,\dots,t_m)=h(t_{\sigma(1)},\dots,t_{\sigma(m)}),$$

for all $\sigma\in S_m$. It follows that terms corresponding to the same permutation in the two equations of Lemma 15 are proportional, with the same constant of proportionality $e^{2c\eta\mu}\epsilon_1$. □

Now we can prove Theorem 13: H_1 has the form $H_1f(\lambda)=\Gamma_1\tilde H_1 f(\lambda)$, where $\tilde H_1$ is the operator of multiplication by

$$\tilde H_1(\lambda)=R_{\Lambda_1,\Lambda_n}(z_1-z_n,\lambda-2\eta\sum_{l=2}^{n-1}h^{(l)})^{(1n)}\cdots R_{\Lambda_1,\Lambda_2}(z_1-z_2,\lambda)^{(12)}. \qquad (7)$$

By using $n-1$ times Lemma 6, we see that

$$\omega(z_2,\dots,z_n,z_1,\lambda)P^{(n,n-1)}\cdots P^{(3,2)}P^{(2,1)}=\omega(z_1,\dots,z_n,\lambda)\tilde H_1(\lambda)^*.$$

Now we use Lemma 16. Let t^* be a solution of (5). Then, for any $u \in V^*_{\Lambda_1}[\mu]$, $v \in (V^*_{\Lambda_2} \otimes \cdots \otimes V^*_{\Lambda_n})[-\mu]$,

$$\omega(z_1, \ldots, z_n, \lambda - 2\eta\mu)\tilde{H}_1(\lambda - 2\eta\mu)^* u \otimes v|_{t^*} = e^{2\eta c\mu}\epsilon_1 \cdot \omega(z_1, \ldots, z_n, \lambda) u \otimes v|_{t^*}. \tag{8}$$

The fact that $\psi(\lambda)$ is an eigenfunction is a consequence of this formula: the function $\psi \in \mathrm{Fun}V[0]$ is uniquely characterized by the property that, for any $w \in (\otimes_j V^*_{\Lambda_j})[0]$, the pairing with $\psi(\lambda)$ is

$$\langle w, \psi(\lambda)\rangle = e^{c\lambda}\omega(z_1, \ldots, z_n, \lambda) w|_{t^*}.$$

Take $w = u \otimes v$, as above. Then $\langle w, H_1\psi(\lambda)\rangle = \langle w, \tilde{H}_1(\lambda - 2\eta\mu)\psi(\lambda - 2\eta\mu)\rangle = e^{c\lambda - 2\eta c\mu}\omega(z_1, \ldots, z_n, \lambda - 2\eta\mu)\tilde{H}_1(\lambda - 2\eta\mu)^* w|_{t^*}$. By (8), this is equal to $\epsilon_1\langle w, \psi(\lambda)\rangle$. Since an arbitrary w can be written as linear combination of vectors of the form $u \otimes v$ as above, the theorem is proved.

5. Completeness of Bethe vectors

We assume that $2\eta = 1/N$, for some positive odd integer N. We also suppose that $\Lambda_1, \ldots, \Lambda_n \in \mathbb{Z}$. This implies that the operators $h^{(i)}$ on $V = V_{\Lambda_1} \otimes \cdots \otimes V_{\Lambda_n}$ have integer eigenvalues. Let $z \in \mathbb{C}^n$ be generic. The commuting difference operators $H_j(z)$ have coefficients which are 1-periodic functions of λ. Thus, for each $\alpha \in \mathbb{C}^\times$ we may consider the eigenvalue problem with multiplier condition

$$\begin{aligned} H_j(z)\psi &= \epsilon_j\psi, \qquad j = 1, \ldots, n, \\ \psi(\lambda + 1) &= \alpha\psi(\lambda). \end{aligned} \tag{9}$$

Let K_N be the field of $1/N$-periodic meromorphic functions of $\lambda \in \mathbb{C}$. The operators $H_j(z)$ are K_N-linear on the $N\dim(V[0])$-dimensional K_N-vector space $\mathcal{H}_\alpha = \{\psi \in \mathrm{Fun}(V[0]) : \psi(\lambda+1) = \alpha\psi(\lambda)\}$.

Theorem 17. *Suppose that $\Lambda_1, \ldots, \Lambda_n$ are integers and let $\sum_{j=1}^n \Lambda = 2m$, $m \in \mathbb{Z}_{\geqslant 0}$. Let $2\eta N = 1$, for some odd integer $N > m$. For generic α, there are $N\dim(V[0])$ solutions of the Bethe ansatz equations (5) with $e^c = (-1)^m\alpha$, such that the corresponding eigenfunctions form a basis of the K_N-vector space $\mathcal{H}_\alpha$*

Proof: Let us first consider the case $m = 1$, and let $t = t_1$ (see the example after Theorem 13). The Bethe ansatz equation is

$$\prod_{l=1}^n \frac{\theta(t - z_l - \eta\Lambda_l)}{\theta(t - z_l + \eta\Lambda_l)} = e^{2c/N},$$

corresponding to the eigenfunction

$$\psi(\lambda) = e^{c\lambda}\sum_{j=1}^n \omega_{(j)}(t, \lambda)\, e_0 \otimes \cdots \otimes e_1 \otimes \cdots \otimes e_0,$$

where $\omega_{(j)}$ stands for $\omega_{0,\ldots,0,1,0,\ldots,0}$ with the 1 at the jth position.

We want to fix the multiplier α. Thus c is of the form $c_0 + 2\pi i r$, $r \in \mathbb{Z}$, where $c_0 = \ln(\alpha) + m\pi i$ for some choice of the branch of the logarithm. Let us now take α real and tending to infinity. Then, for each k, in the vicinity of the point $t_\infty = z_k - \eta\Lambda_k$, we may introduce the local coordinate $u = t - t_\infty$, and the Bethe ansatz equation has the form const $u^{-1}(1 + O(u)) = \alpha^{2/N}e^{-4\pi i s/N}$, for some integer s related to r. This equation has a solution for any sufficiently large α. It has the asymptotic form $t \sim z - a_k + \mathrm{const}\,\alpha^{-2/N}e^{4\pi i s/N}$. Let us denote this solution by $t^{k,s} = t^{k,s}(\alpha)$. The solutions corresponding to different k, s are distinct if the indices run over the sets $k = 1, \ldots, n$,

$s = 0, \dots, N-1$. We claim that the corresponding eigenfunctions $\psi^{k,s}$ are linearly independent over K_N. We use the following simple observation.

Lemma 18. *If $f_1(\lambda), \dots, f_m(\lambda)$ are meromorphic functions on $\mathbb{C}$ with values in $\mathbb{C}^d$ that are linearly dependent over K_N, then the vectors $(f_i(\lambda), f_i(\lambda+1/N), \dots, f_i(\lambda+(N-1)/N))$ in $\mathbb{C}^{md}$ are linearly dependent over $\mathbb{C}$ for all generic $\lambda \in \mathbb{C}$.*

Proof: From $\sum_i a_i(\lambda) f_i(\lambda) = 0$, with a_i $1/N$-periodic, we deduce $\sum_i a_i(\lambda) f_i(\lambda + j/N)$, $j = 0, \dots, N-1$. □

Therefore it is sufficient to show that the vectors

$$\tilde{\psi}^{k,s}(\lambda) = (\psi^{k,s}(\lambda), \psi^{k,s}(\lambda + 1/N), \dots, \psi^{k,s}(\lambda + (N-1)/N)) \in V[0]^N$$

are linearly independent over $\mathbb{C}$ for generic λ.

We prove this by showing that the determinant of these vectors does not vanish for all large enough α. The determinant is proportional to the determinant of the matrix

$$A_{(j,r),(k,s)} = e^{4\pi i s(\lambda + r/N)} \omega_{(j)}(t^{k,s}, \lambda + r/N), \qquad j,k = 1, \dots, n, \qquad r,s = 0, \dots, N-1$$

Let now α tend to infinity. Then the solution $t^{k,s}$ approaches $z_k - a_k$. Thus, $A_{(j,r),(k,s)}$ tends to zero if $k < j$, since $\omega_j(t = z_k - a_k, \lambda) = 0$ for all λ, owing to the vanishing of the factor $\theta(t_j - z_k + a_k)$. Therefore the limiting determinant is block-triangular, and equals the product of determinants of the diagonal N by N blocks. We have

$$\det(A) \to \prod_{j=1}^{n} \det{}_{r,s}(e^{4\pi i s(\lambda + r/N)} \omega_{(j)}(t = z_j - a_j, \lambda + r/N)), \qquad \alpha \to \infty$$

Each factor in the product is, up to multiplication of rows and columns by nonzero factors, the Vandermonde determinant

$$\det{}_{0 \leqslant r,s < N}(e^{4\pi i r s/N}) = \prod_{N > r > s \geqslant 0} (e^{4\pi i r/N} - e^{4\pi i s/N})$$

which does not vanish if N is odd.

The general case is treated along similar lines. We have solutions $t^{K,s}(\alpha)$ labeled by $K = (k_1, \dots, k_n)$ with $k_j \geqslant 0$ and $\sum k_i = m$, and $s \in \{0, \dots, N-1\}$. As the multiplier α tends to infinity, $t^{K,s}$ converges to the point with coordinates

$$t_j = z_l - a_l - 2\eta(k_1 + \cdots + k_l - j), \qquad \text{if } \mathrm{k_1 + \cdots + k_{l-1} < j \leqslant k_1 + \cdots + k_l}.$$

The hypothesis $N > m$ ensures that these coordinates are distinct modulo the lattice. By the same arguments as above, one sees that the linear independence of the corresponding eigenfunctions follows from the non-vanishing of a determinant. This determinant has for $\alpha \to \infty$ a block-triangular form, and the calculation is reduced to the calculation of a Vandermonde determinant. The details are left to the reader. □

Rather than considering the difference operators as acting on the functions of the continuous parameter λ, we may consider the following discrete variant. Let μ be generic and $C_\mu = \{\mu + j/N | j \in \mathbb{Z}\}$. Then the commuting difference operators are defined on the space $\mathrm{Fun}_\mu(V[0])$ of functions of $\lambda \in C_\mu$ with values in $V[0]$, and preserve the subspace $\mathcal{H}_{\alpha,\mu}$ of functions f in $\mathrm{Fun}_\mu(V[0])$ such that $f(\lambda + 1) = \alpha f(\lambda)$.

Corollary 19. *Suppose that $\Lambda_1, \dots, \Lambda_n$ are integers and let $\sum_{j=1}^{n} \Lambda = 2m$, $m \in \mathbb{Z}_{\geqslant 0}$. Let $2\eta N = 1$, for some odd integer $N > m$. For generic α, there are $N \dim(V[0])$*

solutions of the Bethe ansatz equations (5) with $e^c = (-1)^m \alpha$*, such that the corresponding eigenfunctions form a basis of the complex vector space* $\mathcal{H}_{\alpha,\mu}$

It should be possible to treat the case of general rational η (or more generally in $\mathbb{Q}+\tau\mathbb{Q}$) in a similar way.

6. COMPARISON WITH THE ALGEBRAIC BETHE ANSATZ

In this section we compare the Bethe ansatz of this article with the results of [FV2], where eigenfunctions of the transfer matrix of highest weight representations of $E_{\tau,\eta}(sl_2)$ are given.

Let W be a representation of $E_{\tau,\eta}(sl_2)$, see 3.6. Then we have four operators, $a(z,\lambda)$, $b(z,\lambda)$, $c(z,\lambda)$, $d(z,\lambda)$, the matrix elements of the L-operators, acting on W, and obeying the various relations of $E_{\tau,\eta}(sl_2)$. The transfer matrix $T_W(z) \in \mathrm{End}(\mathrm{Fun}(W[0]))$ acts on functions by

$$T_W(z)f(\lambda) = a(z,\lambda)f(\lambda-2\eta) + d(z,\lambda)f(\lambda+2\eta).$$

The relations imply that $T_W(z)T_W(w) = T_W(w)T_W(z)$ for all $z, w \in \mathbb{C}$. If $W = L_1(z_n) \otimes \cdots \otimes L_1(z_1)$, this transfer matrix coincides with the transfer matrix of 2.5 conjugated by

$$\Pi : v_1 \otimes \cdots \otimes v_n \to v_n \otimes \cdots \otimes v_1.$$

Therefore, in this special case, the commuting operators $\Pi H_j(z_1,\dots,z_n)\Pi^{-1}$ are equal to $T_W(z_j)$.

In any case it can be shown in general, using the intertwining property of the R-matrices that $T_W(z)$ *commutes* with the operators $\Pi H_j(z_1,\dots,z_n)\Pi^{-1}$ if $W = V_{\Lambda_n}(z_n) \otimes \cdots \otimes V_{\Lambda_1}(z_1)$.

In [FV2], common eigenfunctions of $T_W(z)$, $z \in \mathbb{C}$ are constructed in the form

$$b(t_1)\cdots b(t_m)v_c$$

where

$$v_c(\lambda) = e^{c\lambda} \prod_{j=1}^{m} \frac{\theta(\lambda - 2\eta j)}{\theta(2\eta)}$$

and $b(t)$ is the difference operator $(b(t)f)(\lambda) = b(t,\lambda)f(\lambda+2\eta)$, $f \in \mathrm{Fun}(W)$ (both the transfer matrix and the difference operators $b(t)$ are part of the *operator algebra* of the elliptic quantum group, see [FV1, FV2]). The variables $t_1,\dots,t_m$ obey a set of Bethe ansatz equations, which are up to a shift the same as the ones described in this paper. The precise relation between the two approaches is the following.

Theorem 20. *Let* $\check{V} = V_{\Lambda_n}(z_n) \otimes \cdots \otimes V_{\Lambda_1}(z_1)$ *be a tensor product of evaluation Verma modules with generic evaluation points* $z_1,\dots,z_n$ *and let*

$$v_c(\lambda) = e^{c\lambda} \prod_{j=1}^{m} \frac{\theta(\lambda - 2\eta j)}{\theta(2\eta)} \in \mathrm{Fun}(\check{V}).$$

Then

$$\begin{aligned}\prod_{j=1}^{m} b(t_j+\eta)\, v_c \;&=\; e^{c(\lambda+2\eta m)}(-1)^m \prod_{i<j} \frac{\theta(t_i - t_j + 2\eta)}{\theta(t_i - t_j)} \\ &\qquad \times \sum_{j_1+\cdots+j_n=m} \omega_{j_1,\dots,j_n}(t_1,\dots,t_n,\lambda)\, e_{j_n} \otimes \cdots \otimes e_{j_1},\end{aligned}$$

where $\omega_{j_1,\dots,j_n}(t_1,\dots,t_n,\lambda)$ *is the image of* $e^*_{j_1} \otimes \cdots \otimes e^*_{j_n}$ *by the map*

$$\omega(z_1,\dots,z_n,\lambda) : V^*_{\Lambda_1} \otimes \cdots \otimes V^*_{\Lambda_n} \to F_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda).$$

Proof: The proof consists of comparing the explicit formula of Proposition 4 with the explicit formula for $\prod b(t_i)\, v_c$ given in [FV2], Theorem 5. $\square$

In particular, this gives an algebraic construction of the spaces $F_a(z,\lambda)$, $a, z \in \mathbb{C}^n$ and their bases: we may define $F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ to be the space spanned by the coordinates of the functions

$$\prod_{i<j} \frac{\theta(t_i - t_j)}{\theta(t_i - t_j + 2\eta)} \prod_{j=1}^{m} b(t_j + \eta)\, v_{c=0}.$$

of $t_1,\dots,t_m$ with values in $\check{V}[\sum \Lambda_j - 2m]$.

From Corollary 19 we thus obtain a completeness result for Bethe vectors of the transfer matrix of the discrete models of [FV2]. Let μ be a generic complex number, and let $C_\mu = \{\mu + j/N | j \in \mathbb{Z}\}$. The transfer matrix $T_{\check{V}}(z)$ of the representation $\check{V}$ is well-defined on the space $\mathrm{Fun}_\mu(\check{V}[0])$ of functions $C_\mu \to \check{V}[0]$ (see [FV2], Section 4).

Theorem 21. *Let $2\eta = 1/N$ for some large enough odd positive integer N. Let $\check{V}$ and v_c be as in Theorem 20, and let $T_{\check{V}}(w)$ be the corresponding transfer matrix acting on functions $C_\mu \to \check{V}[0]$.*

(i) *For any solution $(t_1,\dots,t_m)$ of the Bethe ansatz equations*

$$\prod_{j:j\neq i} \frac{\theta(t_j - t_i - 2\eta)}{\theta(t_j - t_i + 2\eta)} \prod_{k=1}^{n} \frac{\theta(t_i - z_k - (1+\Lambda_k)\eta)}{\theta(t_i - z_k - (1-\Lambda_k)\eta)} = e^{4\eta c}, \qquad i = 1,\dots,m,$$

such that, for all $i<j$, $t_i \neq t_j \mod \mathbb{Z} + \tau\mathbb{Z}$, the vector $\psi = b(t_1)\cdots b(t_m)v \in \mathrm{Fun}_\mu(\check{V}[0])$ is a common eigenvector of all transfer matrices $T_{\check{V}}(w)$ with eigenvalues

$$\epsilon(w) = e^{-2\eta c} \prod_{j=1}^{m} \frac{\theta(t_j - w - 2\eta)}{\theta(t_j - w)} + e^{2\eta c} \prod_{j=1}^{m} \frac{\theta(t_j - w + 2\eta)}{\theta(t_j - w)} \prod_{k=1}^{n} \frac{\theta(w - z_k - (1-\Lambda_k)\eta)}{\theta(w - z_k - (1+\Lambda_k)\eta)}$$

Moreover $\psi(\lambda+1) = (-1)^m e^c \psi(\lambda)$.

(ii) *For any generic $\alpha \in \mathbb{C}^\times$, there are $d = N\dim(\check{V}[0])$ solutions of the Bethe ansatz equations such that the corresponding eigenfunctions form a basis of the space of the d-dimensional vector space of functions $\psi : C_\mu \to \check{V}[0]$ such that $\psi(\lambda+1) = \alpha\psi(\lambda)$.*

Part (i) of this Theorem is taken from [FV1]. Part (ii) follows essentially from Corollary 19. Note, though, that there we had the additional hypothesis that the Λ_i are integers, so that the commuting operators $H_j(z)$ are well-defined on functions on C_μ. However this hypothesis was not used in the proof that the corresponding vectors form a basis. Therefore the proof also applies to the present situation.

Remark. "Large enough" in this theorem means $2N > \Lambda_1 + \cdots + \Lambda_n$, see Theorem 17. This hypothesis can probably be considerably weakened.

Remark. The Bethe ansatz equations have "diagonal" solutions, i.e., solutions for which $t_i = t_j \mod \mathbb{Z} + \tau\mathbb{Z}$ for some $i \neq j$. These solutions do not in general correspond to eigenvectors even if $b(t_1)\cdots b(t_m)v$ is finite and nonzero. The factor

$$\prod_{i<j} \frac{\theta(t_i - t_j)}{\theta(t_i - t_j + 2\eta)}$$

in the correspondence between the eigenvectors obtained by the algebraic Bethe ansatz and the eigenvectors considered in this paper sends the Bethe vectors corresponding to diagonal solutions to zero.

7. Solutions of the qKZB equations

In this section we fix $\tau, \eta, p, \Lambda_1, \dots, \Lambda_n$, and set $a_i = \eta\Lambda_i$.

7.1. Integral representations for solutions.

By a formal Jackson integral solution of the qKZB equations we mean an expression

$$\Psi(z_1, \dots, z_n, \lambda) = \int f(z_1, \dots, z_n, t_1, \dots, t_m, \lambda) Dt_1 \cdots Dt_m,$$

where f takes its values in $V[0]$, which obeys the qKZB equations (1) if we formally use the rule that the "integral" $\int$ is invariant under translations of the variables t_i by p. In other words, $f(z_1, \dots, z_n, t_1, \dots, t_m, \lambda)$ obeys the qKZB equations in the variables z_i up to terms of the form $g(\dots, t_i + p, \dots) - g(\dots, t_i, \dots)$.

Definition: A function $\Phi_a(t)$ depending on a complex parameter a, such that

$$\Phi_a(t+p) = \frac{\theta(t+a)}{\theta(t-a)} \Phi_a(t)$$

is called a (one-variable) phase function.

We assume that p has positive imaginary part, and set $r = e^{2\pi i p}$, $q = e^{2\pi i \tau}$. Then the convergent infinite product

$$\Phi_a(t) = e^{-2\pi i a t/p} \prod_{j=0}^{\infty} \prod_{k=0}^{\infty} \frac{(1 - r^j q^k e^{2\pi i (t-a)})(1 - r^{j+1} q^{k+1} e^{-2\pi i (t+a)})}{(1 - r^j q^k e^{2\pi i (t+a)})(1 - r^{j+1} q^{k+1} e^{-2\pi i (t-a)})}, \tag{10}$$

defines a phase function, and any other phase function is obtained from this by multiplication by a p-periodic function.

Given a one-variable phase function $\Phi_a(t)$, we define with our data an m-variable phase function

$$\Phi(t_1, \dots, t_m, z_1, \dots, z_n) = \prod_{j=1}^{m} \prod_{l=1}^{n} \Phi_{a_l}(t_j - z_l) \prod_{1 \leqslant i < j \leqslant m} \Phi_{-2\eta}(t_i - t_j). \tag{11}$$

Theorem 22. *Let $\Phi_a(t)$ be a phase function, and let Φ be the corresponding m-variable phase function (11). For any entire function ξ of one variable, let*

$$\psi^{\xi}(t, z, \lambda) = \xi(p\lambda - \sum_{l=1}^{n} 2a_l z_l + 4\eta \sum_{j=1}^{m} t_j)) \sum_{j_1 + \cdots + j_n = m} \omega_{j_1, \dots, j_n}(t_1, \dots, t_m, \lambda)\, e_{j_1} \otimes \cdots \otimes e_{j_n}.$$

Then

$$\Psi(z_1, \dots, z_n, \lambda) = \int \Phi(t_1, \dots, t_m, z_1, \dots, z_n) \psi^{\xi}(t_1, \dots, t_m, z_1, \dots, z_n, \lambda) Dt_1 \cdots Dt_m$$

is a formal Jackson integral solution of the qKZB equations.

To obtain solutions from formal Jackson integral solutions, we need to find *cycles*, linear forms on the space of functions of $t_1, \dots, t_m$ that are invariant under translations $t_i \mapsto t_i + p$. To this end we need a stronger version of the preceding theorem, Theorem 24 below, which gives us a space of functions on which our cycles should be defined.

Let Φ be the phase function (11) and let $a = (a_1, \dots, a_n)$, $z = (z_1, \dots, z_n)$. We assume, as usual, that $\sum a_i = 2\eta m$, $m \in \mathbb{Z}_{\geqslant 0}$. For any entire function ξ, let $E^0_a(z; \xi)$, be the space spanned by the functions of $t \in \mathbb{C}^m$ of the form

$$\Phi(t, z) \xi(\lambda - 2\eta \sum_{k=1}^{n} a_k + 4\eta \sum_{j=1}^{m} t_j) f(t, z),$$

where $f(t,z)$, viewed as a function of $t=(t_1,\dots,t_m)$ belongs to $\tilde F^m_a(z,\lambda)$ (see 3.1) for some λ. All components of our integrand belong to this space.

Let $E_a(z;\xi)$, the space of cocycles, be the space spanned by functions of the form $g(t+p\alpha)$, where $g\in E^0_a(z;\xi)$ and $\alpha\in\mathbb{Z}^m$. By construction, $E_a(z;\xi)$ is invariant under translations of the arguments t_i by p. We define the space of coboundaries $DE_a(z;\xi)$ to be the subspace of $E_a(z;\xi)$ spanned by functions of the form $f(\dots,t_j+p,\dots)-f(\dots,t_j,\dots)$, $f\in E_a(z;\xi)$.

Proposition 23. *$E_a(z_1,\dots,z_n)=E_a(z_1,\dots,z_j+p,\dots,z_n)$ for $j=1,\dots,n$*

The proof of this proposition is part of the proof of the following theorem, which is the main result of this section and implies Theorem 22.

Theorem 24. *Let us write the qKZB equations as $\Psi(\dots,z_j+p,\dots)=K_j(z)\Psi(z)$. Then, for all entire functions ξ, the integrand $\Phi(t,z)\psi^\xi(t,z)$ of Theorem 22, viewed as a function of $t\in\mathbb{C}^m$, belongs to $E_a(z;\xi)\otimes V[0]$ for all $z\in\mathbb{C}^n$. It obeys the equations*

$$\Psi(t,\dots,z_j+p,\dots)=K_j(z)\Psi(t,z)\mod DE_a(z;\xi)\otimes V[0],\qquad j=1,\dots,m$$

in the cohomology $(E_a(z;\xi)/DE_a(z;\xi))\otimes V[0]$.

To obtain solutions from these formal solutions, one should find horizontal families of cycles, i.e., linear functions $\gamma(z)$ on $E_a(z;\xi)$ vanishing on $DE_a(z;\xi)$, and such that $\gamma(z+p\alpha)=\gamma(z)$ for all $\alpha\in\mathbb{Z}^n$. This problem will be addressed in the next paper.

7.2. **Proof of Theorem 24.** It is clear that the integrand belongs to $E_a(z;\xi)$. Let us introduce some notation. If f and g are functions of $t_1,\dots,t_m$ such that $\Phi f,\Phi g\in E_a(z;\xi)$, we write $f\sim g$ if $\Phi(t,z)(f(t)-g(t))\in DE_a(z;\xi)$. This means that $f-g$ is a linear combination of expressions of the form Q_ih-h, $\Phi h\in E_a(z;\xi)$, where

$$Q_if(t_1,\dots,t_m)=f(t_1,\dots,t_i+p,\dots,t_m)\phi_i(t)$$

$$\phi_i(t)=\prod_{l=1}^n\frac{\theta(t_i-z_l+a_l)}{\theta(t_i-z_l-a_l)}\prod_{j>i}\frac{\theta(t_i-t_j-2\eta)}{\theta(t_i-t_j+2\eta)}\prod_{j<i}\frac{\theta(t_i-t_j-2\eta+p)}{\theta(t_i-t_j+2\eta+p)}\,,\qquad i=1,\dots,m,$$

The proof of the Theorem is based on the following identity which, similarly to Lemma 16, follows from Lemma 15.

Lemma 25. *Let, for any holomorphic function ξ of one complex variable,*

$$\omega^\xi(z_1,\dots,z_n,\lambda)=\xi(p\lambda-2(\sum_l a_lz_l-2\eta\sum_j t_j))\omega(z_1,\dots,z_n,\lambda).$$

Then, for any $\mu\in\mathbb{C}$ and any $u\in V^*_{\Lambda_1}[\mu]$, $v\in(V^*_{\Lambda_2}\otimes\dots\otimes V^*_{\Lambda_n})[-\mu]$,

$$\prod_{j=1}^m\frac{\theta(t_j-z_1-a_1-p)}{\theta(t_j-z_1+a_1-p)}\omega^\xi(z_1+p,z_2,\dots,z_n,\lambda)u\otimes v\sim\omega^\xi(z_2,\dots,z_m,z_1,\lambda-2\eta\mu)u\otimes v$$

Proof: We first assume that $\xi=1$. We adopt the notation of Lemma 15, and set $\tilde f=\omega(z_1+p,\lambda)$. Then the left-hand side of the claim is, by Lemma 15,

$$\frac{1}{(n-1)!}\mathrm{Sym}\left(\tilde f(t_1,\dots,t_{m'})g(t_{m'+1},\dots,t_m)\prod_{j\leqslant m'}\frac{\theta(t_j-z_1-p+a_1)}{\theta(t_j-z_1-p-a_1)}\right),\tag{12}$$

while the right-hand side is

$$\frac{1}{(n-1)!}\mathrm{Sym}\left(g(t_{m'+1},\dots,t_m)f(t_1,\dots,t_{m'})\prod_{\substack{j\leqslant m'\\ l\geqslant 2}}\frac{\theta(t_j-z_l+a_l)}{\theta(t_j-z_l-a_l)}\prod_{\substack{j\leqslant m'\\ k>m'}}\frac{\theta(t_j-t_k-2\eta)}{\theta(t_j-t_k+2\eta)}\right).$$

To compare these two expressions, notice that $\tilde{f} = \omega(z_1{+}p, \lambda)u$ is related to $f = \omega(z_1, \lambda)u$ by

$$\tilde{f}(t_1 + p, \dots, t_{m'} + p) = f(t_1, \dots, t_{m'}).$$

Therefore we may rewrite the left-hand side in terms of f up to terms which are trivial in the cohomology. For this we use the following formula, valid for general functions h, which is an easy consequence of the definition of Q_i:

$$\begin{aligned} Q_1 \cdots Q_{m'} h(t_1, \dots, t_m) &= h(t_1 + p, \dots, t_{m'} + p, t_{m'+1}, \dots, t_m)\phi^{m'}(t), \\ \phi^{m'}(t) &= \prod_{j=1}^{m'}\prod_{l=1}^{n} \frac{\theta(t_j - z_l + a_l)}{\theta(t_j - z_l - a_l)} \prod_{i \leqslant m' < j} \frac{\theta(t_i - t_j - 2\eta)}{\theta(t_i - t_j + 2\eta)}. \end{aligned}$$

Therefore we have

$$(12) \sim \frac{1}{(n-1)!}\mathrm{Sym}\Big(f(t_1, \dots, t_{m'}) g(t_{m'+1}, \dots, t_m) \prod_{j \leqslant m'} \frac{\theta(t_j - z_1 + a_1)}{\theta(t_j - z_1 - a_1)} \phi^{m'}(t) \Big),$$

which is easily seen to coincide with the right-hand side of the claimed identity.

The case of general ξ is proved in the same way. The point is that the argument of ξ is shifted by $-2a_1 + 4\eta m'$ if $z_1, t_1, \dots, t_{m'}$ are shifted by p, while the same effect is obtained by shifting λ by $-2\eta\mu$. □

Corollary 26. *Let* $1 \leqslant j \leqslant n$ *and let* $w \in V[0]^*$ *be such that* $h^{(j)}w = \mu w$. *Write* $z = (z_1, \dots, z_n)$, $z + p\delta_j = (z_1, \dots, z_j + p, \dots, z_n)$, *and* $\rho_j(t, z) = \prod_{i=1}^{m}[\theta(t_i - z_j + a_j - p)/\theta(t_i - z_j - a_j - p)]$. *Then*

$$\begin{aligned} &\rho_j(t, z)\, \omega^{\xi}(z + p\delta_j, \lambda) R^*_{j-1,j}(z_{j-1} - z_j - p, \lambda - 2\eta \sum_{l<j-1} h^{(l)}) \cdots \\ &\qquad \cdots R^*_{2j}(z_2 - z_j - p, \lambda - 2\eta h^{(1)}) R^*_{1j}(z_1 - z_j - p, \lambda) w \\ &\sim \omega^{\xi}(z_1, \dots, z_n, \lambda - 2\eta\mu) R^*_{j,j+1}(z_j - z_{j+1}, \lambda - 2\eta\mu - 2\eta \sum_{l=1}^{j-1} h^{(l)}) \cdots \\ &\qquad \cdots R^*_{j,n-1}(z_j - z_{n-1}, \lambda - 2\eta \sum_{\substack{l=1 \\ l \neq j}}^{n-2} h^{(l)}) R^*_{jn}(z_j - z_n, \lambda - 2\eta \sum_{\substack{l=1 \\ l \neq j}}^{n-1} h^{(l)}) w \end{aligned}$$

This corollary follows from the above Lemma using Lemma 6.

The proof of Theorem 24 can now be completed: let w be a vector in $V[0]^*$ such that $h^{(j)}w = \mu w$. Let $A(z, \lambda) = R_{1,j}(z_1 - z_j - p, \lambda) \cdots R_{j-1,j}(z_{j-1} - z_j - p, \lambda - 2\eta \sum_{l=1}^{j-2} h^{(l)})$ and $B(z, \lambda) = R_{jn}(z_j - z_n, \lambda - 2\eta \sum_{j \neq l < n} h^{(l)}) \cdots R_{j,j+1}(z_j - z_{j+1}, \lambda - 2\eta \sum_{l<j} h^{(l)})$. With these notations,

$$\begin{aligned} \langle w, A(z, \lambda)\Psi(z{+}p\delta_j, \lambda)\rangle &= \int \Phi(t, z{+}p\delta_j)\, \omega^{\xi}(z{+}p\delta_j, \lambda) A(z, \lambda)^* w \\ &= \int \Phi(t, z)\rho_j(t, z)\, \omega^{\xi}(z{+}p\delta_j, \lambda) A(z, \lambda)^* w \\ &= \int \Phi(t, z)\, \omega^{\xi}(z, \lambda - 2\eta\mu) B(z, \lambda - 2\eta\mu)^* w \\ &= \langle w, B(z, \lambda - 2\eta\mu)\Psi(z, \lambda - 2\eta\mu)\rangle. \end{aligned}$$

The first equality follows from the definition of Ψ. The second is the identity $\Phi(t, z + p\delta_j) = \rho_j(t, z)\Phi(t, z)$, an immediate consequence of the defining property of the phase function. The third equality follows from Corollary 26 and the last equality by the definition of Ψ. Thus $A(z,\lambda)\Psi(z+p\delta_j,\lambda)=\Gamma_j B(z,\lambda)\Psi(z,\lambda)$. To complete the proof of the Theorem, one has to invert the operator A, which is very easy, given the unitarity of R-matrices (property III of Section 2).

8. RESIDUES

In this section we give a proof of Proposition 3, and some other technical results. For this we introduce the basic computational tool of (iterated) residues.

Let us first describe the spaces $F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ in terms of symmetric theta functions. Let $G^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ be the space of meromorphic functions g of m complex variables $t_1,\dots,t_m$ such that

(i) $\prod_{i=1}^m\prod_{k=1}^n\theta(t_i-z_k-a_k)g(t_1,\dots,t_m)$ is a holomorphic function of $t_1,\dots,t_m\in\mathbb{C}$.

(ii) g is periodic with period 1 in each of its arguments and

$$g(\dots,t_j+\tau,\dots)=e^{-2\pi i(\lambda+2\eta m)}g(\dots,t_j,\dots).$$

for all $j=1,\dots,m$.

(iii) $g(t_{\sigma(1)},\dots,t_{\sigma(m)})=g(t_1,\dots,t_m)$ for any permutation $\sigma\in S_m$.

Lemma 27. *The linear map sending f to*

$$g(t_1,\dots,t_m)=\prod_{i<j}\frac{\theta(t_i-t_j+2\eta)}{\theta(t_i-t_j)}f(t_1,\dots,t_m)$$

is an isomorphism from $F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ to $G^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$

Proof: By using the transformation properties of θ under translation by $\mathbb{Z}+\tau\mathbb{Z}$, and the fact that θ is odd, we see that f has the required behavior under lattice translation and the action of permutations if and only if g does. Since the poles of θ lie in the lattice, it is also clear that if g obeys (i), then the poles of f are on the correct hyperplanes. Conversely, if $f\in F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$, then $\prod_{i<j}\theta(t_i-t_j+2\eta)f(t_1,\dots,t_m)$ is regular except possibly on the hyperplanes $t_i\equiv z_j+a_j$ (modulo the lattice). From the symmetry properties of f and the fact that θ is odd, it follows that this function is skew-symmetric under permutation of the arguments. In particular, it vanishes on the diagonals $t_i-t_j=0$, and, by the transformation property of f under translation by $\mathbb{Z}+\tau\mathbb{Z}$, also on the hyperplanes $t_i-t_j=r+s\tau$, r, $s\in\mathbb{Z}$. Therefore we can divide by $\prod_{i<j}\theta(t_i-t_j)$ without creating new poles. Thus g obeys (i). □

Lemma 28. *For any $a_1,\dots,a_n,z_1,\dots,z_n,\lambda\in\mathbb{C}$ and $m\in\mathbb{Z}_{\geqslant 0}$ the dimension of the space $F_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ is*

$$\binom{n+m-1}{m},$$

the number of ways of decomposing m as a sum of n nonnegative integers.

Proof: Multiplication by $\prod_{i=1}^m\prod_{j=1}^n\theta(t_i-z_j-a_j)$ is an isomorphism from the space $G^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ to the space of entire functions that are invariant under permutations of the arguments and have theta function transformation properties under lattice translation of each argument. The dimension of the latter space is easily computed, say by Fourier series. □

In particular, if $n=1$, we have:



FIGURE 1. The point at which the residue is taken.

Lemma 29. *$F_a^m(z,\lambda)$ is a one-dimensional space spanned by*

$$\omega_m(t_1,\dots,t_m) = \prod_{i<j}\frac{\theta(t_i-t_j)}{\theta(t_i-t_j+2\eta)}\prod_{j=1}^{m}\frac{\theta(\lambda+2\eta m+t_j-z-a)}{\theta(t_j-z-a)},$$

The next tool is a suitable set of iterated residues spanning, for generic parameters, the vector space dual to $F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$. For any set $m_1,\dots,m_n$ of nonnegative integers with sum m and any meromorphic function $f(t_1,\dots,t_m)$ in m variables we let $\mathrm{res}_{m_1,\cdots m_n} f$ be the complex number

$$\begin{aligned}\mathrm{res}_{m_1,\dots,m_n} f = {} & \mathrm{res}_{t_1=z_1+a_1-2\eta(m_1-1)}\cdots\mathrm{res}_{t_{m_1-1}=z_1+a_1-2\eta}\mathrm{res}_{t_{m_1}=z_1+a_1}\\ & \mathrm{res}_{t_{m_1+1}=z_2+a_2-2\eta(m_2-1)}\cdots\mathrm{res}_{t_{m_1+m_2-1}=z_2+a_2-2\eta}\mathrm{res}_{t_{m_1+m_2}=z_2+a_2}\\ & \cdots\mathrm{res}_{t_{m-1}=z_n+a_n-2\eta}\mathrm{res}_{t_m=z_n+a_n} f.\end{aligned}$$

The map $f \to \mathrm{res}_{m_1,\dots,m_n} f$ is a linear function on $F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$. The point at which this residue is taken is represented in Fig. 1

Proposition 30. *Fix $a_1,\dots,a_n,z_1,\dots,z_n,\lambda$.*

- (i) *If $z_1,\dots,z_n,\lambda$ and η are generic, then the residues $\mathrm{res}_{m_1,\dots,m_n}$ with $m_1+\cdots+m_n = m$ form a basis of the dual of $F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$.*
- (ii) *Let $\omega_{j_1,\dots,j_n} \in F^m_{a_1,\dots,a_n}(z_1,\dots,z_n,\lambda)$ denotes the image by Φ_n of the basis $\omega_{j_1}\otimes\cdots\otimes\omega_{j_n}$ of $\otimes_{l=1}^n F^{j_l}_{a_l}(z_l,\lambda-2\sum_{k=1}^{l-1}(a_k-2\eta j_k))$, then*

$$\mathrm{res}_{m_1,\dots,m_n}\omega_{j_1,\dots,j_n} = 0$$

  unless $m_l+\cdots+m_n \leqslant j_l+\cdots+j_n$ for all $l = 1,\dots,n$.
- (iii) $\mathrm{res}_{m_1,\dots,m_n}\omega_{m_1,\dots,m_n} \neq 0$ *for generic $z_1,\dots,z_n,\lambda$ and η. More precisely,*

$$\begin{aligned}\mathrm{res}_{m_1,\dots,m_n}\omega_{m_1,\dots,m_n} = {} & \prod_{i=1}^{n}\prod_{j=1}^{m_i}\frac{\theta(\lambda-2\sum_{l<i}(a_l-2\eta m_l)+2\eta j)}{\theta'(0)}\\ & \prod_{1\leqslant l<k\leqslant n}\prod_{j=1}^{m_k}\frac{\theta(z_k+a_k-z_l+a_l-2\eta(j-1))}{\theta(z_k+a_k-z_l-a_l-2\eta(j-1))}\end{aligned}$$

Proof: We prove (ii) and (iii), from which (i) follows.

Let first $n = 1$. By Lemma 28, $F_a(z, \lambda)$ is one dimensional. We compute:

$$\begin{aligned} \mathrm{res}_m \omega_m &= \theta'(0)^{-m} \frac{\prod_{i<j} \theta(2\eta(i-j))}{\prod_{i<j-1} \theta(2\eta(i-j)+2\eta)} \frac{\prod_{j=1}^m \theta(\lambda + 2\eta m - 2\eta(m-j))}{\prod_{j=1}^{m-1} \theta(-2\eta(m-j))} \\ &= \prod_{j=1}^m \frac{\theta(\lambda + 2\eta j)}{\theta'(0)}. \end{aligned}$$

This does not vanish for generic λ.

Consider now the general case.

We introduce the convenient notation $t_I = (t_{i_1}, \dots, t_{i_m})$ for any finite set of integers $I = \{i_1 < \cdots < i_m)$. Then $\omega_{j_1,\dots,j_n}$ can be written as

$$\sum_{I_1,\dots,I_n} \omega_{j_1}(t_{I_1}) \cdots \omega_{j_n}(t_{I_n}) \prod_{l<k} \prod_{i \in I_k} \frac{\theta(t_i - z_l + a_l)}{\theta(t_i - z_l - a_l)} \prod_{i<j, k>l} \prod_{i \in I_k, j \in I_l} \frac{\theta(t_i - t_j - 2\eta)}{\theta(t_i - t_j + 2\eta)}.$$

The sum is taken over all partitions $I_1, \dots, I_n$ of $\{1, \dots, m\}$ such that I_l contains precisely j_l elements, $1 \leqslant l \leqslant m$.

The next step is to determine the partitions corresponding to terms with non-trivial residue $\mathrm{res}_{m_1,\dots,m_n}$. We may assume that all parameters are generic. Then a term with non-trivial residue must in particular have a pole on the hyperplane $t_m = z_n + a_n$, if $m_n > 0$. This implies that $m \in I_n$. If $m_n > 1$, then, we must also have a pole on the hyperplane $t_{m-1} - t_m + 2\eta = 0$. But a pole on the hyperplane $t_j - t_{j+1} + 2\eta$ occurs only if j, $j+1$ belong to the same I_k, (the pole is in ω_k) or if $j \in I_k$, $j+1 \in I_l$ for some $l < k$ (the pole is in the last factor in this case). Thus $m-1 \in I_m$, and so on. We obtain the necessary condition $\{m_1 + \cdots + m_{n-1} + 1, \dots, m\} \subset I_n$. The next residue is at $t_{m_1+\cdots+m_{n-1}} = z_{n-1} + a_{n-1}$. Thus we see that $m_1 + \cdots + m_{n-1}$ must belong to $I_{n-1} \cup I_n$. Continuing this way, we conclude that a partition $I_1, \dots, I_n$ corresponds to a term with trivial residue unless

$$\{m_1 + \cdots + m_l + 1, \dots, m\} \subset I_{l+1} \cup \cdots \cup I_n, \qquad l = 1, \dots, n-1.$$

In particular this proves (ii).

If $m_l = j_l$ for all l, then only one partition contributes to the residue, namely $I_1 = \{1, \dots, m_1\}$, $I_2 = \{m_1 + 1, \dots, m_1 + m_2\}, \dots$ The residue is then the product of the residues of the individual ω_{j_i}'s times the value of the remaining product product of ratios of theta functions, which can easily be computed.

The claim (i) follows from (ii), (iii) and Lemma 28. □

Proof of Proposition 3: By Proposition 30, Φ_n is generically an isomorphism. From this and the associativity of Φ, the main claim of Proposition 3 follows. □

Proof of Lemma 5, (ii): We have to show that $\omega(w, z, \lambda) : V_\Lambda^* \otimes V_\Lambda^* \to F_{aa}(z, w, \lambda)$ is invertible at $z = w$. Since the dimension of $F_{aa}(z, w, \lambda)$ is $m+1$ for all z, w, it is sufficient to show that, for any generic a, the functions $\omega_{j,m-j} = \omega(z, z, \lambda) e_j^* \otimes e_{m-j}^*$, $0 \leqslant j \leqslant m$, are linearly independent. Here we have double poles, the residues $\mathrm{res}_{j,m-j}$ are no longer linearly independent and we have to change slightly the construction. Let $\tilde{\mathrm{res}}_{m,0} = \mathrm{res}_m$ and if $j = 1, \dots, m$, let $\tilde{\mathrm{res}}_{j,m-j} f$ be

$$\begin{gathered} \mathrm{res}_{t_1 = z+a-2\eta(j-1)} \cdots \mathrm{res}_{t_{j-1} = z+a-2\eta} \mathrm{res}_{t_j = z+a} \\ \mathrm{res}_{t_{j+1} = z+a-2\eta(m-j-1)} \cdots \mathrm{res}_{t_{m-1} = z+a-2\eta} \mathrm{res}_{t_m = z+a} [(t_m - z - a) f]. \end{gathered}$$

Then, by proceeding the same way as in the proof of Proposition 30 we see that the matrix $(\tilde{\mathrm{res}}_{j,m-j}\omega_{k,m-k})_{j,k}$ is triangular with non-vanishing diagonal matrix elements. Therefore $\omega(z,z,\lambda)$ is invertible, and

$$\lim_{w\to z} R^*_{\Lambda\Lambda}(z,w,\lambda) = \omega(z,z,\lambda)\omega(z,z,\lambda)^{-1}P = P.$$

□

8.1. **Resonances.** Here we consider the case when the parameters Λ_i are nonnegative integers. From the point of view of elliptic quantum groups (see [FV1]), this is the case when the evaluation Verma module has a finite dimensional quotient. In the present framework, what happens is that the R-matrix has remarkable invariant subspaces.

Let us introduce some notation: if $\ell \in \mathbb{Z}_{\geqslant 0}$, we let $L^*_\ell = \oplus_{j=0}^{\ell}\mathbb{C}e^*_j$, a finite dimensional subspace of V^*_ℓ. Thus its dual space L_ℓ is a quotient of V_ℓ, which we may identify with the subspace $\oplus_{j=0}^{\ell}\mathbb{C}e_j$.

If f is a meromorphic function of m variables $t_1,\dots,t_m$, and $z\in\mathbb{C}$ we define for every integer $\ell\in\{0,\dots,m-1\}$ a function $\mathrm{res}_{(z,\ell)}f$ of $t_1,\dots,t_{m-\ell-1}$:

$$\mathrm{res}_{(z,\ell)}f = \mathrm{res}_{t_{m-\ell}=z-2\eta\ell}\cdots\mathrm{res}_{t_{m-1}=z-2\eta}\mathrm{res}_{t_m=z}.$$

We extend this definition to all $\ell\in\mathbb{Z}_{\geqslant 0}$ by setting $\mathrm{res}_{(z,\ell)}=0$ if $\ell\geqslant m$.

Now consider the map $\omega(z,w,\lambda): V^*_\Lambda\otimes V^*_{\mathrm{M}} \to F_{ab}(z,w,\lambda)$ defined in Section 3. We let $\mathrm{Ker}\,\mathrm{res}_{(u,\ell)}$ be the space of $f\in F_{ab}(z,w,\lambda)$ such that $\mathrm{res}_{(u,\ell)}f=0$.

Theorem 31. *Let z,w,η,λ be generic and $a=\eta\Lambda$, $b=\eta\mathrm{M}\in\mathbb{C}$.*

- (i) *If $\Lambda\in\mathbb{Z}_{\geqslant 0}$, then $\omega(z,w,\lambda)$ maps $L^*_\Lambda\otimes V^*_{\mathrm{M}}$ onto* $\mathrm{Ker}\,\mathrm{res}_{(z+a,\Lambda)}$.
- (ii) *If $\mathrm{M}\in\mathbb{Z}_{\geqslant 0}$, then $\omega(z,w,\lambda)$ maps $V^*_\Lambda\otimes L^*_{\mathrm{M}}$ onto* $\mathrm{Ker}\,\mathrm{res}_{(w+b,\mathrm{M})}$.
- (iii) *If $\Lambda\in\mathbb{Z}_{\geqslant 0}$ and $\mathrm{M}\in\mathbb{Z}_{\geqslant 0}$, then $\omega(z,w,\lambda)$ maps $L^*_\Lambda\otimes L^*_{\mathrm{M}}$ onto* $\mathrm{Ker}\,\mathrm{res}_{(z+a,\Lambda)}\cap\mathrm{Ker}\,\mathrm{res}_{(w+b,\mathrm{M})}$.

Proof: We prove (i). The other claims are proved similarly, or in a simpler way.

Thus we assume that Λ is a nonnegative integer ℓ. We first show that the image of $L^*_\Lambda\otimes V^*_{\mathrm{M}}$ is contained in the kernel of the residue. In other words, we must show that if $f\in F^{m'}_a(z,\lambda)$ and $g\in F^{m''}_b(w,\lambda-2a+4\eta m')$, with $m'\leqslant\ell$, then $\mathrm{res}_{(z+a,\ell)}\Phi(f\otimes g)=0$. Set $m'+m''=m$. The function $\Phi(f\otimes g)$ is, in the notation used in the proof of Proposition 30

$$\sum_{I_1,I_2} f(t_{I_1})g(t_{I_2})\prod_{i\in I_2}\frac{\theta(t_i-z+a)}{\theta(t_i-z-a)}\prod_{i\in I_2,j\in I_1,i<j}\frac{\theta(t_i-t_j-2\eta)}{\theta(t_i-t_j+2\eta)}.$$

The sum runs over all partitions of $\{1,\dots,m\}$ into pairs of disjoint subsets I_1, I_2 of cardinality m', m'', respectively.

The partitions with non-vanishing residue $\mathrm{res}_{(z,\ell)}$ must correspond to terms with poles on the hyperplanes $t_{m-\ell}-t_{m-\ell+1}+2\eta=0,\cdots,t_{m-1}-t_m+2\eta=0$. Thus if $m-1\geqslant j\geqslant m-\ell$ and $j\in I_k$ $(k=1,2)$, then $j+1\in I_r$ with $r\leqslant k$. Since $m'\leqslant\ell$, this implies that $m-\ell\in I_2$. But then the last residue $\mathrm{res}_{t_{m-\ell}=z+a-2\eta\ell}$ vanishes since the pole at $z+a-2\eta\ell=z-a$ is canceled by the zero of $\theta(t_{m-\ell}-z+a)$.

To show that the image is precisely the kernel of the residue, we proceed as in the proof of Proposition 30: suppose $f\in F^m_{ab}(z,w)$ is in Ker $\mathrm{res}_{(z+a,\ell)}$. We claim that f is in the image of $L^*_\Lambda\otimes V^*_{\mathrm{M}}$. Let us think of f as a linear combination of $\omega_{j,m-j}$, $j\in\{0,\dots,m\}$. Consider the residues

$$\widehat{\mathrm{res}}_{m-k,k} = \mathrm{res}_{t_1=w+a-2\eta(m-k-1)}\cdots\mathrm{res}_{t_{m-k}=w+a}\mathrm{res}_{t_{m-k+1}=z+a-2\eta(k-1)}\cdots\mathrm{res}_{t_m=z+a}.$$

As in the proof of Proposition 30, we see that $\widehat{\mathrm{res}}_{m-k,k}\omega_{j,m-j}$ vanishes if $k<j$, and is non-zero for generic z,w,λ, if $k=j$. Also if $f\in\mathrm{Ker}\,\mathrm{res}_{(z+a,\ell)}$, then $\widehat{\mathrm{res}}_{m-k,k}f=0$ whenever

$k > \ell$. In particular, $\widehat{\mathrm{res}}_{m-k,k}\omega_{j,m-j} = 0$ if $j \leqslant \ell$ and $k > \ell$. Therefore, $\widehat{\mathrm{res}}_{m-\ell-1,\ell+1} f = 0$ implies that the coefficient of $\omega_{\ell+1,m-\ell-1}$ vanishes; this and $\widehat{\mathrm{res}}_{m-\ell-2,\ell+2} f = 0$ implies that the coefficient of $\omega_{\ell+2,m-\ell-2}$ vanishes, and so on. We conclude that f is a linear combination of $\omega_{j,m-j}$ with $j \leqslant \ell$, which proves our claim. $\square$

References

[F] G. Felder, *Conformal field theory and integrable systems associated to elliptic curves*, Proceedings of the International Congress of Mathematicians, Zürich 1994, p. 1247–1255, Birkhäuser, 1994; *Elliptic quantum groups,* preprint hep-th/9412207, to appear in the Proceedings of the ICMP, Paris 1994.

[FR] I. Frenkel and N. Reshetikhin, *Quantum affine algebras and holonomic difference equations*, Commun. Math. Phys. 146 (1992), 1–60

[FV1] G. Felder and A. Varchenko, *On representations of the elliptic quantum group $E_{\tau,\eta}(sl_2)$*, preprint q-alg/9601003, to be published in Commun. Math. Phys.

[FV2] G. Felder and A. Varchenko, *Algebraic Bethe ansatz for the elliptic quantum group $E_{\tau,\eta}(sl_2)$*, preprint q-alg/9605024

[T] T. Takebe, *A system of difference equations with elliptic coefficients and Bethe vectors*, preprint q-alg/9604002

[TV1] V. Tarasov and A. Varchenko, *Completeness of Bethe vectors and difference equations with regular singular points*, Int. Math. Res. Notices, (1995), No. 13, 637–669

[TV2] V. Tarasov and A. Varchenko, *Geometry of q-hypergeometric functions as a bridge between Yangians and quantum affine algebras*, preprint q-alg 9604011

[V] A. Varchenko, *Quantized Knizhnik–Zamolodchikov equations, quantum Yang–Baxter equations and difference equations for q-hypergeometric functions*, Commun. Math. Phys. 162 (1994), 499–528